

## Research Note

# The variability analysis of PKS 2155-304

J.H. Fan<sup>1,2</sup> and R.G. Lin<sup>1</sup>

<sup>1</sup> Center for Astrophysics, Guangzhou Normal University, Guangzhou 510400, P.R. China (jhfan@guangztc.edu.cn)

<sup>2</sup> Chinese Academy of Sciences-Peking University Joint Beijing Astrophysical Center (CAS-PKU.BAC), Beijing, P.R. China

Received 9 July 1998 / Accepted 10 September 1999

**Abstract.** In this paper, the post-1977 photometric observations of PKS 2155-304 are compiled and used to discuss the variation periodicity. Largest amplitude variations ( $\Delta U = 1^m.5$ ;  $\Delta B = 1^m.65$ ;  $\Delta V = 1^m.85$ ;  $\Delta R = 1^m.25$ ;  $\Delta I = 1^m.14$ ) and color indices ( $(B-V) = 0.30 \pm 0.06$ ;  $(U-B) = -0.72 \pm 0.08$ ;  $(B-R) = 0.62 \pm 0.07$ ;  $(V-R) = 0.32 \pm 0.04$ ) are found. The Jurkevich's method and DCF (Discrete Correlation Function) method indicate possible periods of 4.16-years and 7.0-years in the V light curve.

**Key words:** galaxies: BL Lacertae objects: individual: PKS 2155-304

## 1. Introduction

BL Lac objects are a special subclass of active galactic nuclei (AGNs) showing some extreme properties: rapid and large variability, high and variable polarization, no or only weak emission lines in its classical definition.

BL Lac objects are variable not only in the optical band, but also in radio, infrared, X-ray, and even  $\gamma$ -ray bands. Some BL Lac objects show that the spectral index changes with the brightness of the source (Bertaud et al. 1973; Brown et al. 1989; Fan 1993), generally, the spectrum flattens when the source brightens, but a different phenomenon has also been found (Fan et al. 1999).

The nature of AGNs is still an open problem; the study of AGN variability can yield valuable information about their nature, and the implications for quasar modeling are extremely important (see Fan et al. 1998a).

PKS 2155-304, the prototype of the X-ray selected BL Lac objects and TeV  $\gamma$ -ray emitter (Chadwick et al. 1999), is one of the brightest and the best studied objects. Its spectrum from  $\lambda 3600$  to  $\lambda 6800$  appears blue ( $B-V < 0.1$ ) and featureless (Wade et al. 1979). A 0.17 redshift was claimed from the potentially detected weak [O III] emission feature (Charles et al. 1979), which was not detected in Miller & McAlister (1983) observations. Later, a redshift of 0.117 was obtained from several discrete absorption features (Bowyer et al. 1984). PKS 2155-304 varies at all observation frequencies and is one of the most

extensively studied objects for both space-based observations in UV and X-ray bands (Treves et al. 1989; Urry et al. 1993; Pian et al. 1996; Giommi et al. 1998) and multiwavelength observations (Pesce et al. 1997). Variation over a time scale of one day was observed (Miller & Carini 1991) and that over a time scale of as short as 15 minutes is also reported by Paltani et al. (1997) in the optical band. Different brightness-dependent spectrum properties are found (see Miller & McAlister 1983; Smith & Sitko 1991; Urry et al. 1993; Courvoisier et al. 1995; Xie et al. 1996; Zhang & Xie 1996; Paltani et al. 1997).

In this paper, we will investigate the periodicity in the light curve and discuss the variation as well. The paper has been arranged as follows: In Sect. 2, the variations are presented and the periodicities are searched, in Sect. 3, some discussion and a brief conclusion are given.


## 2. Variation

### 2.1. Light curves

The optical data used here are from the literature: Brindle et al. (1986); Carini & Miller (1992); Courvoisier et al. (1995); Griffiths et al. (1979); Hamuy & Maza (1987); Jannuzi et al. (1993); Mead et al. (1990); Miller & McAlister (1983); Pesce et al. (1997); Smith & Sitko (1991); Treves et al. (1989); Urry et al. (1993); Xie et al. (1996) and shown in Fig. 1a-e. From the data, the largest amplitude variabilities in UBVRI bands are found:  $\Delta U = 1^m.5(11^m.87 - 13^m.37)$ ;  $\Delta B = 1^m.65(12^m.55 - 14^m.20)$ ;  $\Delta V = 1^m.85(12^m.27 - 14^m.13)$ ;  $\Delta R = 1^m.25(11^m.96 - 13^m.21)$ ;  $\Delta I = 1^m.14(11^m.55 - 12^m.69)$  and color indexes are found:  $(B-V) = 0.30 \pm 0.06$  (N=140 pairs);  $(U-B) = -0.72 \pm 0.08$  (N=105 pairs);  $(B-R) = 0.62 \pm 0.07$  (N=90 pairs);  $(V-R) = 0.32 \pm 0.04$  (N=98 pairs), the uncertainty is  $1\sigma$  dispersion.

### 2.2. Periodicity

The photometric observations of PKS 2155-304 indicate that it is variable on time scales ranging from days to years (Miller & McAlister 1983). Is there any periodicity in the light curve? To answer this question, the Jurkevich (1971) method is used



**Fig. 1.** a: The long-term U light curve of PKS 2155-304; b: The long-term B light curve of PKS 2155-304; c: The long-term V light curve of PKS 2155-304; d: The long-term R light curve of PKS 2155-304; e: The long-term I light curve of PKS 2155-304.

to search for the periodicity in the V light curve since there are more observations in this band.

The Jurkevich method (Jurkevich 1971, also see Fan et al. 1998a) is based on the expected mean square deviation and it is less inclined to generate spurious periodicity than the Fourier analysis. It tests a run of trial periods around which the data are folded. All data are assigned to  $m$  groups according to their phases around each trial period. The variance  $V_i^2$  for each group and the sum  $V_m^2$  of all groups are computed. If a trial period equals the true one, then  $V_m^2$  reaches its minimum. So, a “good” period will give a much reduced variance relative to those given by other false trial periods and with almost constant values. To show the significance of the trial periodicity, we adopted the  $F$ -test (see Press et al. 1992).

When the Jerkevich method is used to V measurements, some results are obtained and shown in Fig. 2 ( $m = 10$ ), which shows several minima corresponding to trial periods of less than 4.0-years and two broad minima corresponding to averaged periods of  $(4.16 \pm 0.2)$  and  $(7.0 \pm 0.16)$  years respectively.


For the periods, which are smaller than 4.0-years, we found that the decrease of the  $V_m^2$  is less than 3 times the noise suggest-

ing that it is difficult for one to take them as real signatures of periods, i.e., those periods should be discussed with more observations. For the two broad minima, the  $F$ -test is used to check their reality. The significance level is 93.8% for the 4.16-year period and 96.2% for the 7.0-year period.


### 3. Discussion

PKS 2155-304 was observed more than 100 years ago. Griffiths et al. (1979) constructed the annually averaged B light curve up to the 1950’s from Harvard photographic collection. But there are only a few observations during the period of 1950-1970. The periodicity obtained here (see Fig. 2) are based on the post-1977 data.

For comparison, we adopted the DCF (Discrete Correlation Function) method to the V measurements. The DCF method, described in detail by Edelson & Krolik (1988) (also see Fan et al. 1998b), is intended for analyses of the correlation of two data sets. This method can indicate the correlation of two variable temporal series with a time lag, and can be applied to the periodicity analysis of a unique temporal data set. If there is a period,  $P$ , in the lightcurve, then the DCF should show clearly



**Fig. 2.** Plot of  $V_m^2$  vs. trial period,  $P$ , in years.



**Fig. 3.** DCF for the V band data. It shows that the V light curve is self-correlated with time lags of 4.2 and 7.31 years. In addition, there are also correlation with time lags of less than 4.0 years.

whether the data set is correlated with the DCF with time lags of  $\tau = 0$  and  $\tau = P$ . It can be done as follows.

Firstly, we have calculated the set of unbinned correlation (UDCF) between data points in the two data streams  $a$  and  $b$ , i.e.

$$UDCF_{ij} = \frac{(a_i - \bar{a}) \times (b_j - \bar{b})}{\sqrt{\sigma_a^2 \times \sigma_b^2}}, \quad (1)$$

where  $a_i$  and  $b_j$  are points in the data sets,  $\bar{a}$  and  $\bar{b}$  are the average values of the data sets, and  $\sigma_a$  and  $\sigma_b$  are the corresponding standard deviations. Secondly, we have averaged the points sharing


the same time lag by binning the  $UDCF_{ij}$  in suitably sized time-bins in order to get the  $DCF$  for each time lag  $\tau$ :

$$DCF(\tau) = \frac{1}{M} \sum UDCF_{ij}(\tau), \quad (2)$$

where  $M$  is the total number of pairs. The standard error for each bin is

$$\sigma(\tau) = \frac{1}{M-1} \{\sum [UDCF_{ij} - DCF(\tau)]^2\}^{0.5}. \quad (3)$$

The resulting DCF is shown in Fig. 3. Correlations are found with time lags of  $(4.20 \pm 0.2)$  and  $(7.31 \pm 0.16)$  years. In ad-



**Fig. 4.** The observed V light curve (filled points) and the simulated V light curve (solid curve) with the periods of 4.16 and 7.0 considered.

dition, there are signatures of correlation with time lags of less than 3.0 years. If we consider the two minima in both the right and left sides of the 7.0-year minimum, then we can say that the periods of 4.16 and 7.0-years found with the Jerkevich method are consistent with the time lags of 4.2-years and 7.3-years found with the DCF method. These two periods are used to simulate the light curve (see the solid curve in Fig. 4).

It is clear that the solid curve does not fit the observations so well. One of the reasons is that there are probably more than two periods ( $\sim 4.2$  and  $\sim 7.0$  years) in the light curve as the results in Fig. 2 and 3 indicate. Another reason is that the derived period is not so significant as Press (1978) mentioned. Press argued that periods in the order of one third of the time span have a high probability of appearing if longer-term variations exist. The data used here have a time coverage of about 16.0 years, i.e., about 3 times the derived periods. Therefore, these are only tentative and should be confirmed by independent work.

From the data, the largest amplitude variations are found for UBVR bands with I and R bands showing smaller amplitude variations. One of the reasons is from the fact that there are fewer observations for those two bands, another reason is perhaps from the effect of the host galaxy, which affects the two bands more seriously.

In this paper, the post-1970 UBVR data are compiled for 2155-304 to discuss the spectral index properties and to search for the periodicity. Possible periods of 4.16 and 7.0 years are found.

*Acknowledgements.* We are grateful to the referee for his/her comments and suggestions! This work is supported by the National Pan Deng Project of P.R. China and the National Natural Scientific Foundation of P.R. China and the National 973 project of P.R. China.

## References

- Bertaud C.H., Wlerick G., Véron P., et al. 1973, *A&AS* 11, 77  
 Bowyer S., Brodie J., Clarke J. T., Henry J.P. 1984, *ApJ* 278, L103  
 Brindle C., Hough J.H., Bailey J.A., et al. 1986, *MNRAS* 221, 739  
 Brown L.M.J., Robson E.I., Gear W.K., Smith M.G.: 1989, *ApJ* 340, 150  
 Chadwick P.M., Lyons K., McComb T.J.L., et al. 1999, *ApJ* 513, 161  
 Carini M.T., Miller H.R., 1992, *ApJ* 385, 146  
 Charles P., Thorstensen J., Bowyer S., 1979, *Nat.* 281, 285  
 Courvoisier T.J.L., Blecha A., Bouchet P., et al., 1995, *ApJ* 438, 108  
 Edelson, R.A. & Krolik, J.H., 1988, *ApJ*, 333, 646  
 Fan J.H., Xie G.Z., Pecontal E., et al., 1998a, *ApJ*, 507, 173  
 Fan J.H., Adam G. Xie G.Z. et al., 1998b, *A&AS*, 136, 217  
 Fan J.H.: 1993, Ph.D. Thesis, Yunnan Astronomic Observatory, Chinese Academy of Sciences.  
 Fan J.H., Xie G.Z., Adam G., et al. 1999, *ASP Conf. Ser.* Vol. 159, Eds. L. Takalo & A. Sillanpaa, p99  
 Giommi P., Fiore F., Guainazzi M., et al. 1998, *A&A* 333, L5  
 Griffiths R. E., Briel U., Chaisson L., Tapia S., 1979, *ApJ* 234, 546  
 Hamuy M., Maza J., 1987, *A&AS* 68, 383  
 Jurkevich I. 1971, *Ap&SS* 13, 154  
 Jannuzi B.T., Smith P.S., Elston R., 1993, *ApJS* 85, 265  
 Mead A.R.G., Ballard K.R., Brand P.W.J.L., et al. 1990, *A&AS* 83, 183  
 Miller H.R., Carini M.T., 1991, in *Variability of Active Galactic Nuclei*, eds. H.R. Miller and P.J. Wiita (Cambridge: Cambridge Univ. Press), p256  
 Miller H.R., McAlister H.A., 1983, *ApJ* 272, 26  
 Paltani S., Courvoisier T.J.L., Blecha A., Bratschi P., 1997, *A&A* 327 539  
 Pesce J.E., Urry C.M., Maraschi L., et al. 1997, *ApJ* 486, 770  
 Pian E., Urry C.M., Pesce J.E. et al. 1996, in "Blazar Continuum Variability" eds. H.R. Miller, J.R. Webb, and J.C. Noble, *ASP Conf. Series* Vol. 110, 417

- Press W.H. 1978 *Comm. on Astrophys.* 7, 103
- Press W.H., Teukolsky S.A., Vetterling W.T., Flannery B.P. 1992, in *Numerical Recipes in C, The Art of Scientific Computing* (2nd Edition), Cambridge Uni. Press.
- Smith P.S., Sitko M.L., 1991, *ApJ* 383, 580
- Treves A., Morini M., Chiappetti L. et al. 1989, *ApJ* 341, 733
- Urry C.M., Maraschi L., Edelson R. et al. 1993, *ApJ* 411, 614
- Wade R.A., Szkody P., Córdova F., 1979, *IAU Circ.*, No. 3279.
- Zhang Y.H., Xie G.Z. 1996, *A&AS* 119, 199
- Xie G.Z., Zhang Y.H., Li K.H. et al. 1996, *AJ*, 111, 1065