

Mid-infrared ISO spectroscopy of NGC 4945*

H.W.W. Spoon¹, J. Koornneef², A.F.M. Moorwood¹, D. Lutz³, and A.G.G.M. Tielens²

¹ European Southern Observatory, Karl-Schwarzschild-Strasse 2, 85748 Garching, Germany (hspoon, amoor@eso.org)

² Kapteyn Astronomical Institute, Postbus 800, 9700 AV Groningen, The Netherlands (j.koornneef, tielens@astro.rug.nl)

³ Max-Planck-Institut für Extraterrestrische Physik (MPE), Postfach 1603, 85740 Garching, Germany (lutz@mpe.mpg.de)

Received 28 January 2000 / Accepted 20 March 2000

Abstract. We have observed the central region of the nearby starburst galaxy NGC 4945 with the mid-infrared spectrometers SWS and PHT-S aboard ISO. We do not find any evidence for the existence of the powerful AGN, inferred from hard X-ray observations. The upper limits on our AGN tracers [Ne V] 14.32 μm & 24.3 μm and [Ne VI] 7.65 μm imply an $A_V > 160$ towards the NLR, assuming the NLR to be of equal strength as in the Circinus galaxy. Other possibilities are discussed. The starburst excitation indicators [Ne III] 15.56 μm / [Ne II] 12.81 μm and $L_{\text{bol}}/L_{\text{lyc}}$ suggest that the starburst in the central region is at least 5×10^6 yrs old, and that it accounts for at least half of the nuclear bolometric luminosity. The starburst might well power the entire bolometric luminosity, but the available constraints are also consistent with an up to 50% contribution of the embedded AGN. With PHT-S, at a resolution of ≈ 90 , we detect strong absorption features of water ice, and, for the first time in an external galaxy, of CO₂ and CO. The same PHT-S spectrum also reveals strong emission from the family of PAH features. Finally, we have observed and detected several pure rotational and ro-vibrational H₂ lines, two of which, the (0-0) S(0) & S(1) lines, allow us to determine the excitation temperature (160 K) and warm H₂ mass ($2.4 \times 10^7 M_{\odot}$). The low excitation temperature shows Orion-like shocks not to be representative for the entire emission of the central region of the galaxy and fairly normal PDRs to be perhaps more typical.

Key words: galaxies: individual: NGC 4945 – galaxies: ISM – galaxies: starburst – galaxies: Seyfert

1. Introduction

NGC 4945 is a nearby, large (20' \times 4') spiral galaxy seen nearly edge on ($i \sim 78^\circ$; Ott 1995). At a recession velocity of 560 km s^{-1} it is at the mean radial velocity of the Centaurus group (Hesser et al. 1984), of which it is believed to be a member. Distance estimates vary between 3.5 and 4.0 Mpc (see Bergman

et al. 1992 and Mauersberger et al. 1996 for discussions). In this paper we will adopt a distance of 3.9 Mpc (Bergman et al. 1992), which implies that 1'' is equivalent to 18 pc.

NGC 4945 is one of the brightest infrared galaxies in the sky: S[12]=24 Jy, S[25]=43 Jy, S[60]=588 Jy, S[100]=1416 Jy (Rice et al. 1988). The total infrared luminosity amounts to $L(8\text{--}1000 \mu\text{m}) = 2.95 \times 10^{10} L_{\odot}$, $\sim 75\%$ of which originates in the central $12'' \times 9''$ (Brock et al. 1988).

Near infrared observations reveal the nuclear region to be the site of a powerful, yet visually obscured, starburst. Br γ (Moorwood et al. 1996a) and Pa α (Marconi et al. 2000) recombination line maps show the starburst to be concentrated in a circumnuclear disk or ring ~ 200 pc across (11''). Further evidence for (a period of) strong star formation comes from the discovery of a conical structure, roughly perpendicular to the galaxy major axis. It is believed to be a cavity, vacated by a starburst-driven superwind (Heckman et al. 1990; Moorwood et al. 1996a). The non-detection of [O III] within the cone and the absence of coronal lines excludes an AGN as the driver of the outflow.

Clear evidence for the presence of an AGN comes from hard X-ray observations (Iwasawa et al. 1993; Guainazzi et al. 2000). The AGN X-ray emission is however heavily absorbed by a column density of $10^{24.7} \text{ cm}^{-2}$, which obscures the AGN at all optical and infrared wavelengths. Previous authors have attributed most of the IR luminosity to the starburst (e.g. Moorwood & Oliva 1994; Koornneef & Israel 1996). Hard X-ray observations with *BeppoSAX* indicate that the bolometric luminosity may as well be accounted for by the AGN alone (Guainazzi et al. 2000).

3 cm & 6 cm ATCA radio maps of the central region of NGC 4945 (Forbes & Norris 1998) are dominated by strong nuclear emission, and emission extended along the disc of the galaxy. There is also evidence for some filamentary structure associated with the cavity cleared by the starburst superwind. VLBI observations by Sadler et al. (1995) reveal the existence of a compact radio core. This, as well as the presence of H₂O megamasers in a Keplerian disc about a $\sim 10^6 M_{\odot}$ black hole (Greenhill et al. 1997), are taken as further evidence for the presence of an AGN.

Near infrared observations of molecular hydrogen emission in NGC 4945 have been reported by several authors over the last 15 years (e.g. Moorwood & Glass 1984; Moorwood & Oliva

Send offprint requests to: H.W.W. Spoon

* Based on observations with ISO, an ESA project with instruments funded by ESA Member States (especially the PI countries: France, Germany, the Netherlands and the United Kingdom) and with the participation of ISAS and NASA

1988; Koornneef 1993; Moorwood & Oliva 1994; Koornneef & Israel 1996; Moorwood et al. 1996a; Quillen et al. 1999; Marconi et al. 2000). While fluxes are known for eight rovibrational transitions accessible from the ground (Koornneef & Israel 1996), spatial information is available only for the (1-0) S(1) 2.1218 μm line. These observations show the H₂ emission to be associated with the hollow cone, *not* with the starburst traced in hydrogen recombination emission. The absence of a correlation argues against photons as the source of excitation. Instead, the emission is attributed to shock heating of the molecular material at the face of the cavity (Moorwood et al. 1996a; Marconi et al. 2000).

Mid-infrared spectroscopy is much less affected by intervening extinction than the UV and optical equivalents, with $A(\lambda)/A_V$ less than 0.1. Observations of the central region of the galaxy, using the mid-infrared spectrometer SWS (De Graauw et al. 1996) and the spectrophotometer PHT-S (Lemke et al. 1996), both aboard ISO (Kessler et al. 1996), are therefore very useful to study the nuclear components otherwise hidden by heavy extinction. In Sect. 3.1 we present the results of the search for high excitation emission from the AGN. In Sect. 3.2 we study the properties of the nuclear starburst. In Sect. 3.3 we discuss the dominant nuclear power source. Sect. 3.4 discusses the broad emission and absorption features, tracing the properties of the interstellar medium in and in front of the nucleus. Finally, in Sect. 3.5 we discuss the physical conditions and excitation of the warm molecular hydrogen.

2. Observations

As part of the Central Program “MPEXGAL”, we have observed the central region of NGC 4945 with the Short Wavelength Spectrometer (SWS) and the spectrophotometer PHT-S on board ISO.

2.1. SWS spectroscopy

SWS grating line profile scans (SWS02 mode) were obtained on 1996 February 6 for 28 spectral lines in the range of 2.42 to 40.34 μm . The spectral resolution in this range varies between $R=900$ and 2000, corresponding to a velocity resolution of 330–150 km s^{-1} . Aperture sizes used range between $14'' \times 20''$ and $20'' \times 33''$. SWS was centered on the 1.4GHz continuum peak from Ables et al. (1987), which coincides with the the position of the L-band peak of Moorwood et al. (1996a). At the time of observation the position angle of the major axis of the SWS apertures was 35.4° , 10° off from that of the galaxy major axis (45°).

The data reduction was performed using the SWS Interactive Analysis system (SIA; Lahuis et al. 1998; Wieprecht et al. 1998), putting special emphasis on tools to improve cosmic ray spike removal, dark current subtraction and flat fielding. The wavelength calibration of SWS is discussed by Valentijn et al. (1996a). We used calibration files as of March 1999. The accuracy of the flux calibration is estimated to be 30% (Schaeidt et al. 1996). The resulting spectra are shown in Fig 1.

Table 1. NGC 4945 results from SWS observations

Identification	λ_{rest} [μm]	F_{obs} [W/cm^2]	Aperture [$'' \times ''$]
H ₂ (1-0) Q(3)	2.424	$3.20 \cdot 10^{-20}$	14×20
H I Br α	4.052	$7.79 \cdot 10^{-20}$	14×20
H ₂ (0-0) S(7)	5.510	$1.11 \cdot 10^{-19}$	14×20
H ₂ (0-0) S(5)	6.909	$1.54 \cdot 10^{-19}$	14×20
H I Pf α	7.460	$1.95 \cdot 10^{-20}$	14×20
[Ne VI]	7.652	$<7.3 \cdot 10^{-21}$	14×20
[Ar III]	8.991	$<2.3 \cdot 10^{-20}$	14×20
[S IV]	10.511	$<8.8 \cdot 10^{-21}$	14×20
H ₂ (0-0) S(2)	12.279	$7.45 \cdot 10^{-20}$	14×27
[Ne II]	12.814	$8.68 \cdot 10^{-19}$	14×27
[Ne V]	14.322	$<7.0 \cdot 10^{-21}$	14×27
[Cl II]	14.368	$5.19 \cdot 10^{-21}$	14×27
[Ne III]	15.555	$7.51 \cdot 10^{-20}$	14×27
H ₂ (0-0) S(1)	17.035	$1.51 \cdot 10^{-19}$	14×27
[S III]	18.713	$6.88 \cdot 10^{-20}$	14×27
[Ar III]	21.829	$<5.1 \cdot 10^{-21}$	14×27
[Fe III]	22.925	$6.46 \cdot 10^{-21}$	14×27
[Fe I]	24.042	$<5.4 \cdot 10^{-21}$	14×27
[Ne V]	24.318	$<5.5 \cdot 10^{-21}$	14×27
[S I]	25.257	$<1.4 \cdot 10^{-20}$	14×27
[O IV]	25.890	$3.00 \cdot 10^{-20}$	14×27
[Fe II]	25.988	$4.11 \cdot 10^{-20}$	14×27
H ₂ (0-0) S(0)	28.221	$4.82 \cdot 10^{-20}$	20×27
[S III]	33.481	$4.87 \cdot 10^{-19}$	20×33
[Si II]	34.815	$9.66 \cdot 10^{-19}$	20×33
[Fe II]	35.349	$<2.8 \cdot 10^{-20}$	20×33
[Ne III]	36.014	$<1.6 \cdot 10^{-20}$	20×33
o-H ₂ O	40.341	$<2.8 \cdot 10^{-20}$	20×33

In total 17 spectral lines were detected. For another 11 lines we derived upper limits, using gaussian profiles of width equal to other lines of the same (or comparable) species, scaled to a peak height corresponding to approximately 3σ of the noise. Both detections and upper limits are presented in Table 1.

2.2. PHT-S spectrophotometry

We have obtained two low resolution ($\lambda/\Delta\lambda \sim 90$) ISO-PHT-S spectra, on 1996 October 12 and 1997 August 3, respectively. ISO-PHT-S comprises two low-resolution grating spectrometers covering simultaneously the wavelength range 2.47 to 4.87 μm and 5.84 to 11.62 μm . The spectrum is registered by two linear arrays of 64 Si:Ga detectors with a common entrance aperture of $24'' \times 24''$. The measurements were carried out in rectangular chopped mode, using a chopper throw of $180''$. The resulting spectra thus are free of contributions from zodiacal light, that would otherwise affect the spectrum. The pure on-source integration times were 512 and 1024 s.

The ISO-PHT-S data were reduced using PIA¹ (Gabriel et al. 1997) version 8.1. Steps in the data reduction included: 1) deglitching on ramp level. 2) subdivision of ramps in two sec-

¹ PIA is a joint development by the ESA Astrophysics Division and the ISO-PHT Consortium




Fig. 1. SWS line spectra of NGC 4945. Typical $\pm 1\sigma$ error bars are marked. Note that noise increases towards the edges of scans

tions of 32 non destructive read-outs. 3) ramp fitting to derive signals. 4) masking of bad signals by eye-inspection. 5) kappa sigma and min/max clipping on remaining signal distribution. 6) determination of average signal per chopper plateau. 7) masking or correction of bad plateaux by eye-inspection. 8) background subtraction using all but the first four plateaux. 9) finally, flux calibration, using the signal dependent spectral response function of Acosta-Pulido (1999). This spectral response function avoids some deficiencies of the previous PIA response function, in particular in the $3 \mu\text{m}$ region near the “Ice” feature. The absolute calibration is accurate to within 20%.

The two resulting spectra were obtained at slightly different position angles about the nucleus. For the first, the square aperture was aligned with the galaxy major axis (45°). For the second, the position angle was 31.1° . Fig. 2 shows the averaged ISO-PHT-S spectrum. The on-source integration times

were used as weight factors in the computation of the average spectrum.

A number of emission lines can be identified in the ISO-PHT-S spectrum. These include $9.66 \mu\text{m}$ H_2 (0-0) S(3), the unresolved blend of $6.99 \mu\text{m}$ [Ar II] and $6.91 \mu\text{m}$ H_2 (0-0) S(5), and $4.05 \mu\text{m}$ H Br α . For the $9.66 \mu\text{m}$ H_2 (0-0) S(3) line, not included in the SWS02 line scans, we measure a flux of $5.4 \times 10^{-20} \text{ W/cm}^2$, with an uncertainty of 30%.

3. Results

3.1. AGN not seen at mid-infrared wavelengths

NGC 4945 is a peculiar and interesting target for studying the relation of AGN and star formation in galaxies. Clear evidence for hidden AGN activity comes from hard X-ray observations. NGC 4945 is amongst the brightest hard X-ray emitting galaxies




Fig. 2. The average ISO-PHT-S spectrum of NGC 4945. Upper inset: comparison with the ISO-SWS spectrum of the line of sight towards the Galactic center. Lower inset: the ISO-PHT-SL spectrum in linear flux scale

and exhibits variability of its 13–200 keV flux on timescales of ~ 10 hrs, which clearly establishes its AGN origin (Iwasawa et al. 1993; Guainazzi et al. 2000; Marconi et al. 2000). The AGN X-ray emission is heavily absorbed by a column density of $10^{24.7} \text{ cm}^{-2}$ (corresponding to $A_V \sim 2600$), a high value, but within the range observed for Seyfert 2 galaxies (e.g. Risaliti et al. 1999). In unified schemes, the X-ray obscuration measures a line of sight towards the very center. Obscuration towards the NLR probes a different line of sight and is usually significantly lower, making the NLR visible in Seyfert 2 galaxies.

Mid-infrared high excitation lines are able to penetrate a far larger dust obscuration than their optical and UV counterparts. They are therefore ideally suited as tracers of embedded AGN activity. Mid-infrared emission lines like [Ne V] $14.32 \mu\text{m}$ & $24.32 \mu\text{m}$, [Ne VI] $7.65 \mu\text{m}$ and [O IV] $25.9 \mu\text{m}$ are prominently present in the spectrum of all Seyferts observed with ISO (Moorwood et al. 1996b; Genzel et al. 1998). On the other hand, the same emission lines are also weakly visible towards, for instance, supernova remnant RCW 103 (Oliva et al. 1999). [O IV] emission (at the few percent level compared to [Ne II]) has also been detected in a sample of starburst galaxies (Lutz et al. 1998), again at a level much weaker than seen in typical AGNs. The origin of the weak level emission in these sources is believed to be shocks. A detection of any of the high excitation lines discussed above does therefore not automatically imply the detection of an AGN in NGC 4945.

We do not detect the lines of [Ne V] and [Ne VI]. No trace of [Ne VI] is seen in the wing of the nearby PAH emission feature (Fig. 1). From Fig. 1 it might appear that the two [Ne V] lines were indeed detected. However, at the level at which the features appear, instrumental effects play a significant role. In the $14.35 \mu\text{m}$ line scan a strong fringe in the relative spectral response function coincides *exactly* with the expected position for the [Ne V] line. Depending on the size of the emitting area, the feature may be entirely attributed to this instrumental effect. We therefore chose to state an upper limit for the $14.32 \mu\text{m}$

[Ne V] line. The feature seen in the other [Ne V] scan, centered at $24.4 \mu\text{m}$, was registered by only two detectors, although 12 detectors scanned over the central wavelength. As visible in Fig. 1, the feature is redshifted with respect to the NGC 4945 systemic velocity. This shift is not observed for any other line we observed. We therefore derive an upper limit for this [Ne V] line too.

The only detected high ionization line in NGC 4945 is the $25.9 \mu\text{m}$ [O IV] line. An AGN contribution to this line is possible – to match the limits on higher excitation lines, only part of the [O IV] emission would be related to an AGN. The detection of possibly shock-related [O IV] in many starbursts (Lutz et al. 1998) cautions, however, that this may be a more plausible origin of [O IV] in NGC 4945. The ratio of 0.033 with respect to $[\text{Ne II}] + 0.44 \times [\text{Ne III}]$ is above average, but well within the range observed for the Lutz et al. (1998) starbursts, also considering that the high extinction (Sect. 3.2) will increase the observed ratio relative to the intrinsic one. A population of Wolf-Rayet stars as origin of the [O IV] emission seems unlikely. Lutz et al. (1998) have shown that the [O IV] emission would have to originate in widely dispersed small H II regions and would have to be relatively strong. [O IV] emission at this level has not been observed in local star forming regions. A conservative analysis will hence not attribute the [O IV] emission in NGC 4945 to the AGN nor to a population of Wolf-Rayet stars.

The limits on high excitation AGN tracers are consistent with several scenarios, or perhaps more likely a combination of them:

- **The Narrow Line Region is extremely obscured even in the mid-IR.** We derive an $A_V \geq 160$ ($A(7.65 \mu\text{m}) = A(14.3 \mu\text{m}) = A(24.3 \mu\text{m}) \geq 4.3$) to the NLR from a comparison of Circinus and NGC 4945 [Ne V] and [Ne VI] line strengths, under the assumption that the galaxies' NLRs are similar. The choice for Circinus is motivated in Table 2.

Table 2. In X-ray properties NGC 4945 is very similar to Circinus. However, IR indicators for a narrow line region are missing in NGC 4945

	NGC 4945	Circinus
Distance [Mpc]	3.9	4 ^a
H col. dens. to AGN [cm ⁻²]	5 × 10 ²⁴	4 × 10 ^{24b}
A _V to AGN	2600	2100
L(2–10 keV) [L _⊙]	78 × 10 ^{7c}	8.8–44 × 10 ^{7b}
νL _ν (100 keV)/L _{FIR}	0.003 ^d	0.002 ^d
L(8–1000 μm) [L _⊙]	2.2 × 10 ¹⁰	1.2 × 10 ^{10a}
[Ne v]14.3 μm [W/cm ²]	<7.0 × 10 ⁻²¹	440 × 10 ^{-21e}
[Ne v]24.3 μm [W/cm ²]	<5.5 × 10 ⁻²¹	244 × 10 ^{-21e}
[Ne VI]7.65 μm [W/cm ²]	<7.3 × 10 ⁻²¹	413 × 10 ^{-21e}

^a Siebenmorgen et al. (1997)

^b Matt et al. (1999)

^c Guainazzi et al. (2000)

^d Marconi et al. (2000)

^e Moorwood et al. (1996b)

- **UV photons from the AGN are absorbed close to the nucleus along all lines of sight**
- **The extreme ultraviolet luminosity of the AGN is lower than in Circinus.** In comparison to the Circinus SED, this would imply a large deficiency in UV relative to X-ray flux (Table 2).

3.2. Starburst properties

Near-infrared broad-band and emission-line imaging has revealed the nucleus of NGC 4945 to be the site of a sizeable starburst, the presence of which is illustrated by the conically shaped starburst superwind-blown cavity traced at many near-infrared wavelengths (Moorwood et al. 1996a; Marconi et al. 2000). Hampered by the large extinction even in the near-infrared, age estimates for the nuclear starburst are sparse and intrinsically uncertain. ISO-SWS offers the possibility for the first time to observe the mid-infrared line ratio [Ne III] 15.56 μm/[Ne II] 12.81 μm. This ratio, which is much less affected by extinction than visible and UV lines, is sensitive to the hardness of the stellar radiation field and hence is a good indicator for the age of the nuclear starburst. We observed the two lines in the same ISO-SWS aperture, which was centered on the nucleus (see Table 1).

To estimate the extinction to the NGC 4945 nuclear starburst we use the ratio of the 18.71 μm and 33.48 μm [S III] lines. This ratio is commonly used as a density diagnostic for the density range 10^{2.5}–10^{4.5} cm⁻³ and is only mildly dependent on the temperature of the emitting gas. Assuming a typical starburst gas density of 300 cm⁻³ (Kunze et al. 1996; Rigopoulou et al. 1996), the intrinsic ratio should be ~0.43 (i.e. the value in the low density limit, computed using the collision strengths of Tayal & Gupta 1999). The observed ratio is far lower: 0.14 ± 0.06. We hence deduce a screen extinction of A(18.7 μm) = 1.7^{+0.8}_{-0.5}, which, using the Galactic center extinction law of Draine (1989; with A(9.7 μm)/E(J–K) = 0.71),

amounts to A_V = 36⁺¹⁸₋₁₁. This value is to be considered an upper limit in case the [S III] emitting area is larger than 14'' × 27'', in which case an aperture effect causes the [S III] ratio to be a lower limit.

Another independent estimate of the extinction is usually obtained from hydrogen recombination line strengths, assuming ‘case-B’ conditions. For NGC 4945 we therefore observed the 4.05 μm Brα and the 7.46 μm Pfα line. Both were measured in the same aperture. The ratio Pfα/Brα is 0.25 ± 0.10, whereas ‘case-B’ recombination theory predicts a ratio of 0.32. The extinction at 7.46 μm must therefore be similar or slightly *larger* than at 4.05 μm. This indicates that the grain composition is unusual and probably more similar to the composition found in the line of sight towards the Galactic center (Lutz et al. 1996; Lutz 1999) than found towards other parts of our Galaxy. An extinction towards the NGC 4945 nuclear starburst can therefore at present not be derived from lines in the 4–7 μm range.

The extinction derived for the nuclear starburst is somewhat larger than the value we derive for the warm molecular hydrogen (see Sect. 3.5). This indicates that the warm molecular hydrogen and nuclear starburst emission are coming from different nuclear components, the latter possibly more enshrouded than the former. With the unusual grain composition in mind, it is striking how well the Galactic Center extinction law fits our molecular hydrogen data, resulting in a smooth excitation diagram, even for the H₂ (0-0) S(3) line in the center of the 9.7 μm silicate feature (see Sect. 3.5). We are therefore confident that the extinction correction for the starburst, derived using the [S III] ratio, is also useful.

In order to determine the excitation of the nuclear starburst we apply the extinction correction derived from the [S III] ratio to the observed [Ne III]/[Ne II] ratio. The extinction corrections amount to A(12.8 μm) = 1.51 and A(15.6 μm) = 1.19. The extinction corrected [Ne III]/[Ne II] ratio is 0.064^{+0.037}_{-0.032}. Thornley et al. (2000) list observed [Ne III]/[Ne II] ratios for 26 starburst galaxies, all measured in the same ISO-SWS configuration. Clearly, NGC 4945 is among the lowest excitation targets in their sample. Note that the ISO-SWS aperture used is large compared to the typical size scales in starbursts. The ratios listed by Thornley et al. (2000) should therefore be regarded as aperture averaged.

For starburst galaxies L_{bol}/L_{lyc} is another measure of the excitation of star clusters. Depending on the upper mass cut-off, the star formation decay time scale and the age of the clusters, Thornley et al. (2000) modeled L_{bol}/L_{lyc} to lie between 3 and 200. The measured values for starburst galaxies range between 3 and 50. Below we will determine L_{bol}/L_{lyc} for the NGC 4945 nuclear starburst. We assume L_{bol} = L_{IR} (i.e. *no* AGN contribution to L_{IR}) and estimate L_{lyc} from the dereddened 4.05 μm Brα flux. For A(4.05 μm) = 1.2 (applying the Galactic center law of Draine (1989) for A_V = 36⁺¹⁸₋₁₁) and L_{lyc} = 670 L_{Brα} we find L_{lyc} = 8⁺⁹₋₄ × 10⁸ L_⊙ and L_{bol}/L_{lyc} = 28⁺²⁶₋₁₅. Using the 12.81 μm [Ne II] line and the empirical scaling L_{lyc} = 64 L_{Ne II} (Genzel et al. 1998) a similar result is obtained.

Given the variety of possible star forming histories, it is hard to constrain the age of the nuclear starburst (assuming *no* AGN contribution to L_{IR}). However, both excitation diagnostics agree

on a low excitation which suggests an evolved burst with an age in excess of 5×10^6 years, but would also be consistent with a low IMF upper mass cut-off.

Marconi et al. (2000) show that it is possible to construct starburst models which are consistent with their near-infrared observations of NGC 4945, but differ by the total luminosity generated (their Fig. 4). An instantaneous burst would not leave space in the energy budget for a sizable contribution from the hidden AGN, whereas a combination of instantaneous burst and constant star formation would. We would like to point out here that the latter model would be inconsistent with the low $[\text{Ne III}]/[\text{Ne II}]$ ratio observed by us. Only their instantaneous burst is in agreement with both the near-infrared and mid-infrared observations.

3.3. What powers the nucleus of NGC 4945?

The large extinction towards the nuclear starburst and the AGN buried within, makes it very difficult to assess the contributions of either component to the nuclear bolometric luminosity. The optical, near-infrared, mid-infrared and far-infrared spectra of NGC 4945 are entirely consistent with a starburst-like nature: BLR or NLR high-excitation lines are absent; the starburst excitation indicator $[\text{Ne III}]/[\text{Ne II}]$ has a starburst-like value; the ratios of $6 \mu\text{m}$ (ISO-PHT-S), S12 or S25 to S60 or S100 fluxes are all very low and consistent with emission from cold dust only. Furthermore, the ratio $L_{\text{bol}}/L_{\text{lyc}} = 28^{+26}_{-15}$, is perfectly consistent with the low excitation of the starburst as deduced from $[\text{Ne III}]/[\text{Ne II}]$. And last, NGC 4945 lies on the radio-far-infrared correlation for starburst galaxies (Forbes & Norris 1998). Hence, the starburst might well account for the the entire observed bolometric luminosity.

On the other hand, Guainazzi et al. (2000), who have observed the AGN in NGC 4945 in hard X-rays, compute the AGN to be able to account for all the bolometric luminosity observed, *if* it has a typical quasar L_X/L_{bol} ratio. Since there is no such thing as a template AGN spectrum, the conversion factor applied, $L_{1-10 \text{ keV}}/L_{\text{bol}} \sim 0.05$ (Elvis et al. 1994), may have an uncertainty which could easily allow for the NGC 4945 starburst to dominate the bolometric luminosity instead.

The same uncertainties surround the accretion rate of the $\sim 1.6 \times 10^6 M_{\odot}$ black hole inferred from H_2O maser observations (Greenhill et al. 1997). A high but not implausible rate of 50% of the Eddington rate ($L_{\text{Edd}} \sim 4.1 \times 10^{10} L_{\odot}$) would suffice to power the observed bolometric luminosity. Given the wide range of efficiencies inferred for Seyferts, this information does not add anything to identify the dominant power source.

In this complex situation with two potentially dominant power sources, the most significant constraint on their relative weight is the total $L_{\text{bol}}/L_{\text{lyc}}$ ratio and its implications on the $L_{\text{bol}}/L_{\text{lyc}}$ of the *starburst* component. $L_{\text{lyc}}^{\text{sb}}$ is directly constrained by observations, but $L_{\text{bol}}^{\text{sb}}$ changes for different assumptions on the starburst and AGN contributions to the total bolometric luminosity. If there is a significant AGN contribution, $(L_{\text{bol}}/L_{\text{lyc}})_{\text{sb}}$ will be less than the global value of 28. Values as low as ~ 3 which are possible for a zero age massive star pop-

ulation with Salpeter IMF (e.g. Leitherer & Heckman 1995) are strongly inconsistent with the low excitation observed in NGC 4945. Thornley et al. (2000) model $[\text{Ne III}]/[\text{Ne II}]$ and $L_{\text{bol}}/L_{\text{lyc}}$ ratios of starbursts, taking into account clusters of different ages within the ISO-SWS aperture. An evolving starburst with $[\text{Ne III}]/[\text{Ne II}] = 0.064$ as in NGC 4945 must have a $L_{\text{bol}}/L_{\text{lyc}} \geq 15$ (their Fig. 8). This limit simply reflects the higher $L_{\text{bol}}/L_{\text{lyc}}$ of later type O stars and persists if the low excitation is due to an upper mass cut-off rather than evolution. With a lower limit of ~ 15 on $(L_{\text{bol}}/L_{\text{lyc}})_{\text{sb}}$, the starburst contribution to the bolometric luminosity must be at least $\sim 50\%$.

We hence conclude that the AGN in NGC 4945 plays a secondary although most likely not insignificant role in the energetics of this nearby galaxy. Extremely small values for the AGN contribution to the bolometric luminosity would imply an unrealistically high ratio of L_X/L_{bol} for the AGN. The very low inferred black hole mass, the very cold mid-infrared to far-infrared colors, and the absence of any clear line of sight towards the AGN, support our view that starburst activity dominates AGN activity in NGC 4945.

3.4. Emission and absorption features

The infrared spectrum of the central region of NGC 4945 obtained with ISO-PHT-S (see Fig. 2) presents a new view of the ISM in starburst galaxies. Even at the low spectral resolving power of $R \approx 90$ the spectrum is dominated by a wealth of emission and absorption features.

Especially prominent is the family of PAH emission features at 3.3, 6.2, 7.7, 8.6 and $11.3 \mu\text{m}$, which ISO confirmed to be common-place in most galactic and extragalactic ISM spectra (e.g. Acosta-Pulido et al. 1996; Rigopoulou et al. 1999; Mattila et al. 1999; Clavel et al. 2000). Nevertheless, the weakness of the 8.6 and $11.3 \mu\text{m}$ PAH bands in NGC 4945 is unusual. Consistent with $A_V \sim 36$ and with the strength of the absorption features discussed below, we explain this weakness by heavy extinction, which will suppress these two features placed in the wings of the silicate absorption feature.

Perhaps the most important result, however, is the rich absorption spectrum, indicating that we are observing the infrared sources in the central region of NGC 4945 through a medium containing molecular ices. Interstellar absorptions of $4.27 \mu\text{m}$ (2343 cm^{-1}) solid CO_2 and $4.68\text{--}4.67 \mu\text{m}$ 'XCN'+CO are detected, the first time in an extragalactic source to our knowledge. At our resolving power and signal-to-noise we cannot determine the contribution of $4.62 \mu\text{m}$ (2165 cm^{-1}) XCN, $4.67 \mu\text{m}$ (2140 cm^{-1}) CO ice and of gaseous CO absorptions to the $4.58\text{--}4.67 \mu\text{m}$ absorption complex. The strength of the XCN absorption appears to be remarkable, suggestive of ice grain processing in an energetic environment (Lacy et al. 1984). We defer a more detailed analysis of the XCN/CO feature to a future paper, which will also include the results of follow-up observations with ISAAC at the VLT. A strong silicate feature is observed around $9.7 \mu\text{m}$ (see also Moorwood & Glass 1984). A deep minimum is also detected around $3.0 \mu\text{m}$, which is suggestive of water ice (or more precise, the O-H stretch) absorption. Ta-

ble 3 gives column densities for some of the absorption features discussed above. The presence and strength of these absorption features is consistent with the high starburst obscuration derived from the emission lines (but see also Chiar et al. (2000) for variations in the strength of features along lines of sight of similar A_V).

At the resolution of ISO-PHT-S the molecular absorption features in NGC 4945 show striking similarities with the features seen in the ISO-SWS spectrum of the line of sight towards the Galactic center (Lutz et al. 1996; see Fig. 2). Observations at our spectral resolution do however not permit a detailed comparison. Regarding the $4.26 \mu\text{m}$ CO_2 feature it is likely that the feature can be attributed to solid state CO_2 , since high spectral resolution ISO-SWS observations of other sources indicate that the contribution of gaseous CO_2 to the observed feature is very small (see in particular van Dishoeck et al. 1996). In the $4.6\text{--}4.8 \mu\text{m}$ region, the spectra of NGC 4945 and the Galactic center differ more strongly, and a more careful inspection is required to assess the contributions of gaseous and solid CO and XCN. ISO-SWS spectroscopy of the Galactic center (Lutz et al. 1996) clearly shows that what we see at low resolution as a relatively shallow and broad feature is in fact dominated by individual lines of gaseous CO. Contributions of a potential underlying solid CO/XCN component are possible but difficult to separate until our high resolution follow-up observations have been executed.

3.5. Molecular hydrogen:

physical conditions, excitation and mass

Near infrared observations of molecular hydrogen emission in NGC 4945 have been reported by several authors over the last 15 years. The most complete set of observations is published by Koornneef & Israel (1996), who observed 8 ro-vibrational transitions with IRSPEC at the ESO NTT. With ISO-SWS and ISO-PHT-S we have extended the number of observed lines from 8 to 14 by observing the pure rotational transitions S(0), S(1), S(2), S(3), S(5) and S(7) as well as the (1-0) Q(3) line. The latter was also observed with IRSPEC and can therefore be used to determine the proper aperture correction factor for the IRSPEC data set. An overview of the observed lines is presented in Table 4.

Information on the spatial extent of the H_2 emitting region is only available for the $2.12 \mu\text{m}$ (1-0) S(1) line (Koornneef & Israel 1996; Moorwood et al. 1996a; Quillen et al. 1999; Marconi et al. 2000). Based on Fig. 3a of Moorwood et al. (1996a) we estimate that more than 90% of the (1-0) S(1) emission fits within the smallest ISO-SWS aperture ($14'' \times 20''$). It is not unreasonable to expect the H_2 emitting area to increase with decreasing H_2 temperature. The aperture sizes used to observe the respective H_2 transitions increase with increasing sensitivity to lower temperature H_2 . Based on this, we will assume in what follows that ISO-SWS and ISO-PHT-S have observed all available warm H_2 . Further to this, all three instruments were centered on the same nuclear position and viewed the nuclear region under more or less similar position angles (see Sect. 2).

We will use the ratio of the 1-0 Q(3) line fluxes measured by ISO-SWS and IRSPEC to scale the other IRSPEC lines to the ISO-SWS aperture size. This ratio is 2.33.

From the 14 transitions observed it is possible to compute the upper level populations for 12 levels. We assumed the H_2 levels to be optically thin. The excitation diagram in Fig. 3 shows a plot of the natural logarithm of the total number of H_2 molecules (N_u), divided by the statistical weight (g_u), in the upper level of each transition detected, versus the energy of that level (E_u/k). The plot shows the situation after extinction correction (see below).

The excitation temperature (T_{ex}) of the gas is the reciprocal of the slope of the excitation diagram. If the warm H_2 is in LTE, the excitation temperature directly corresponds to the kinetic temperature. As is clearly visible from Fig. 3, (extinction corrected; see below) the excitation temperature increases monotonically with upper level energy, from 160 K for the combination of (0-0) S(0) & S(1) to 2200 K for the ro-vibrational lines.

In a highly obscured galaxy like NGC 4945, extinction corrections to the H_2 emission will be important. The extinction can be estimated from the H_2 data themselves taking into account that any known excitation mechanism should produce a “smooth” excitation diagram for the pure rotational lines, and that transitions originating in a common upper level should give consistent results. More specifically, we use three criteria:

- The excitation temperature should increase monotonically from the lowest to the highest energy levels. This sets limits on the extinction correction for the (0-0) S(3) line in the center of the $9.7 \mu\text{m}$ silicate absorption feature.
- The ratio of the (1-0) Q(3) & (1-0) S(1) lines at 2.42 & $2.12 \mu\text{m}$ should be its intrinsic ratio determined by molecular constants only. The same applies to the (1-0) Q(2) & (1-0) S(0) lines at 2.41 & $2.22 \mu\text{m}$, that originate from identical upper levels too.
- In LTE, the upper level populations normalized by the statistical weights should be similar for the 0-0 S(7) & 1-0 Q(3) lines at 5.51 & $2.42 \mu\text{m}$, which differ by only 4% in upper level energy.

We have varied the extinction and tried several extinction laws. We present the most applicable extinction laws here:

- Law A: $A(\lambda) \propto \lambda^{-1.75}$ for $\lambda < 8 \mu\text{m}$. For $\lambda > 8 \mu\text{m}$ we took the Galactic center law of Draine (1989), with $A(9.7 \mu\text{m})/E(J-K)=0.71$ (Roche & Aitken 1985) and $E(J-K)=5$.
- Law B: The same as law ‘A’, except for the range $\lambda=[2.6, 8] \mu\text{m}$, where we took the extinction law as found towards the Galactic center (Lutz 1999). In the $3\text{--}8 \mu\text{m}$ range this reddening law constitutes a significantly higher extinction than usually assumed.

From Fig. 3 and the criteria defined above, moderate extinctions of $A_V=17\text{--}23$ are clearly preferred. Extinction law A provides a somewhat better fit than extinction law B. None of the 3 solutions gives a good fit to the (1-0) Q(4) data point.

Table 3. Column densities of solid state features towards the nucleus of NGC 4945 and towards the Galactic center (SgrA*). In order to derive the column densities we integrated $\int \tau_\nu d\nu$ over the width of the band and divided the result by the bandstrength A. N_{H} was determined from $N_{\text{H}}=1.9 \times 10^{21} A_V$, where $A_V=30$ for SgrA* and $A_V=36$ for NGC 4945

Species	λ_{rest} [μm]	A [cm/mol.]	τ_{center}		N [mol./cm ²]		N/N _H	
			NGC 4945	SgrA*	NGC 4945	SgrA*	NGC 4945	SgrA*
H ₂ O	3.09	$2 \times 10^{-16\text{a}}$	0.9	0.50 ^b	2.4×10^{18}	$1.3 \times 10^{18\text{b}}$	3.5×10^{-5}	2.3×10^{-5}
CO ₂	4.27	$7.4 \times 10^{-17\text{c}}$	0.8 ^d	0.72 ^e	2.0×10^{17}	$1.7 \times 10^{17\text{e}}$	2.9×10^{-6}	3.0×10^{-6}
XCN	4.60	–	–	–	–	–	–	–

^a Hagen & Tielens (1981)

^b Chiar et al. (2000)

^c Gerakines et al. (1995)

^d Determined by fitting a Gaussian profile followed by rebinning to the ISO-PHT-S instrument resolution

^e Gerakines et al. (1999)

Table 4. NGC 4945 molecular hydrogen data. $A(\lambda)$ is the extinction correction in magnitudes; A_{ul} is the Einstein coefficient for the transition from level u to level l. T_u is the upper level energy of level u; g_u is the statistical weight of level u; $N_u(\nu, J)$ is the number of H₂ molecules in upper level u

Identification	λ_{rest} [μm]	$F_{\text{obs}}^{\text{a}}$ [W/cm ²]	Instrument	Aperture ["' × "']	$A(\lambda)^{\text{b}}$	A_{ul} [1/s]	T_u [K]	g_u	$N_u(\nu, J)/g_u^{\text{c}}$
0-0 S(0)	28.2207	4.82×10^{-20}	SWS	20×27	0.44	2.94×10^{-11}	510	5	1.27×10^{62}
0-0 S(1)	17.0346	1.51×10^{-19}	SWS	14×27	0.84	4.76×10^{-10}	1015	21	5.13×10^{60}
0-0 S(2)	12.2785	7.45×10^{-20}	SWS	14×27	1.04	2.76×10^{-9}	1682	9	8.79×10^{59}
0-0 S(3)	9.6649	5.43×10^{-20}	PHT-S	24×24	2.34	9.84×10^{-9}	2504	33	1.28×10^{59}
0-0 S(5)	6.9091	1.54×10^{-19}	SWS	14×20	0.27	5.88×10^{-8}	4587	45	4.74×10^{57}
1-0 Q(1)	2.4066	1.48×10^{-20}	IRSPEC	6×6	1.72	4.29×10^{-7}	6150	9	9.69×10^{56}
1-0 Q(2)	2.4134	5.1×10^{-21}	IRSPEC	6×6	1.71	3.03×10^{-7}	6471	5	8.47×10^{56}
1-0 S(0)	2.2235	4.4×10^{-21}	IRSPEC	6×6	1.98	2.53×10^{-7}	6471	5	1.03×10^{57}
1-0 Q(3)	2.4237	3.20×10^{-20}	SWS	14×20	1.70	2.78×10^{-7}	6951	21	5.82×10^{56}
1-0 Q(3)	2.4237	1.36×10^{-20}	IRSPEC	6×6	1.70	2.78×10^{-7}	6951	21	5.82×10^{56}
1-0 S(1)	2.1218	1.29×10^{-20}	IRSPEC	6×6	2.15	3.47×10^{-7}	6951	21	5.84×10^{56}
0-0 S(7)	5.5103	1.11×10^{-19}	SWS	14×20	0.40	2.00×10^{-7}	7197	57	7.13×10^{56}
1-0 Q(4)	2.4375	2.2×10^{-21}	IRSPEC	6×6	1.68	2.65×10^{-7}	7584	9	2.28×10^{56}
2-1 S(1)	2.2477	2.3×10^{-21}	IRSPEC	6×6	1.94	4.98×10^{-7}	12550	21	6.36×10^{55}
2-1 S(2)	2.1542	0.6×10^{-21}	IRSPEC	6×6	2.09	5.60×10^{-7}	13150	9	3.79×10^{55}

^a Before aperture correction

^b Extinction law 'A' & $A_V=20$ (see text)

^c After aperture correction (IRSPEC data only) and extinction correction. Adopted distance $D=3.9\text{Mpc}$




Fig. 3. Excitation diagram for molecular hydrogen in NGC 4945. Different symbols are used to distinguish different instrumental origin: *diamond*: ISO-SWS; *square*: ISO-PHT-S; and *cross*: IRSPEC. Results are shown for three different dereddening schemes, marked by different shades of grey. *Light-grey* denotes the combination extinction law 'B' & $A_V=50$, *middle-grey* denotes extinction law 'B' & $A_V=20$ and *black* extinction law 'A' & $A_V=20$

Table 5. Warm molecular hydrogen mass estimates for the nucleus of NGC 4945 using the best fit power law $dM/dT=4.43 \times 10^{15} T^{-4.793} M_{\odot}/K$. The total H_2 gas mass estimated from CO amounts to $2.7 \times 10^8 M_{\odot}$.

Temp. range [K]	M(warm H_2) [M_{\odot}]	% of total M(H_2)
200–10000	2.19×10^6	0.8%
150–10000	6.51×10^6	2.4%
120–10000	1.52×10^7	5.6%
100–10000	3.03×10^7	11%
90–10000	4.52×10^7	17%
80–10000	7.06×10^7	26%
70–10000	1.17×10^8	43%
60–10000	2.10×10^8	78%
50–10000	4.20×10^8	156%

In the following analysis, we use the preferred extinction correction of $A_V=20_{-3}^{+3}$ and extinction law A. We note that the extinction to the H_2 emitting region is slightly less than that to the starburst [H II] regions (Sect. 3.2). This plausibly matches the morphological results of Moorwood et al. (1996a), who find the starburst in an obscured disk, but the H_2 emission extending into a less obscured wind blown cavity.

A rough estimate of the amount of warm molecular hydrogen in the nucleus of NGC 4945 can be derived from the level populations of the pure rotational S(0) and S(1) transitions. The excitation temperature for the upper levels of these transitions (J=2 and J=3) is 160 K. Assuming the same excitation conditions for the J=0 and J=1 levels, we compute a warm molecular hydrogen mass of $2.4 \times 10^7 M_{\odot}$. This is 9% of the total H_2 gas mass estimated from CO and 0.7% of the dynamical mass interior to the molecular ring (Bergman et al. 1992; see below).

As already noted, the excitation temperature changes significantly with level energy. This is the consequence of the natural fact that the emitting gas will consist of a mixture of temperatures. The rich NGC 4945 dataset allows us to address this in a more quantitative way. Experiments with fits assuming a number of discrete temperature components lead us to suggest that a power law might give a good representation of the mass distribution as a function of temperature. We obtain a good fit for the following power law: $dM/dT=4.43 \times 10^{15} T^{-4.793} M_{\odot}/K$. The quality of the fit is shown in Fig. 4.

Table 5 gives warm molecular masses for several low temperature cut-offs. Since H_2 gas at temperatures below 70 K does not contribute to the (0-0) S(0) flux, nor to any of the other pure rotational lines, we cannot verify whether our power law mass distribution continues down to the lowest temperatures. Nevertheless, we have included mass estimates down to a low temperature cut-off of 50 K. This number is reasonable, since we don't expect the giant molecular clouds (GMCs) to be as cold as in the Galactic disk (10–20 K). Rather we expect conditions as found near the Galactic center, where the GMCs are believed to have temperatures exceeding 50 K (Armstrong & Barrett 1985).

It is interesting to compare our warm molecular hydrogen gas mass estimate with values found in the literature (see

Moorwood & Oliva 1994 for a review). Bergman et al. (1992), using the inner molecular rotation curve of Whiteoak et al. (1990), compute a dynamical mass interior to the molecular ring ($R \leq 280 \text{ pc} = 15.6''$) of $3.3 \times 10^9 M_{\odot}$. The same authors use CO to derive a total H_2 gas mass of $2.7 \times 10^8 M_{\odot}$ for the ring, assuming the rather high kinetic gas temperature of 100 K. Note that a low temperature cut-off of the order 50–60 K in our H_2 temperature distribution would bring our estimate of the total H_2 gas mass in agreement with that derived from the low level CO observations. The total H_2 gas mass of $2.7 \times 10^8 M_{\odot}$ agrees well with a starburst-like position of NGC 4945 in the $L_{\text{IR}}-M(H_2)$ diagram (Young & Scoville 1991).

In Table 6 we list for a number of external galaxies and Galactic template sources temperatures and masses of the warm molecular hydrogen gas. With a value of 9%, NGC 4945 has a warm H_2 gas fraction similar to that found for most of the other galaxies listed. Note however that the results for NGC 3256, NGC 4038/39 and Arp 220 are less well constrained than for NGC 6946 and NGC 4945: only for the latter two can the temperature of the warm H_2 gas be determined from the (0-0) S(0) and S(1) transitions directly. For the same reason a comparison of the H_2 gas temperatures is of limited value unless they are derived from identical line combinations. Fairly low temperatures can be derived from the (0-0) S(0) and S(1) lines, 160 K and 179 K for NGC 4945 and NGC 6946, respectively. Limits for other galaxies listed in Table 6 are consistent with a similarly low temperature. This temperature is well below that observed for an Orion type shock (>500 K). It is closer to what is observed for the same line combination in PDRs (e.g. Orion Bar: 155 K, D. Rosenthal priv. comm.; S140: 159 K, Draine & Bertoldi, 1999). While a variety of regions may contribute to the galaxy-integrated temperature distribution, this comparison clearly shows Orion-like shocks to be not representative for the entire emission, and fairly normal PDRs (or less energetic shocks) to be perhaps more typical. If excited by shocks (as suggested by the morphology, Moorwood et al. 1996a), then the near-infrared H_2 emission in NGC 4945 must trace a small subcomponent of faster shocks.

4. Conclusions

The main results of this paper can be summarized as follows:

- The nuclear starburst is heavily obscured by 36_{-11}^{+18} mag. of visual extinction, as inferred from the [S III] $18.7 \mu\text{m}/33.5 \mu\text{m}$ ratio.
- The excitation of the nuclear starburst is very low, as deduced from excitation indicators [Ne III] $15.56 \mu\text{m}/$ [Ne II] $12.81 \mu\text{m}$ and $L_{\text{bol}}/L_{\text{lyc}}$, consistent with an age of at least 5×10^6 yrs. Comparison with starburst models implies that at least 50% of the bolometric luminosity is powered by the starburst.
- The very low inferred black hole mass, the very cold mid-infrared to far-infrared colors, and the absence of any free line of sight to the NLR supports the conclusion that the starburst dominates the bolometric luminosity.




Fig. 4. Ratio of the fitted and observed H₂ excitation diagram for the best fitting power law $dM/dT=4.43 \times 10^{15} T^{-4.793} M_{\odot}/K$

Table 6. Warm molecular hydrogen in external galaxies and Galactic template sources. In column 4 detected lines are printed bold, upper limits normal. T_{01} refers to the excitation temperature computed from the (0-0) S(0) & S(1) fluxes. Likewise, T_{12} refers to the excitation temperature computed from the (0-0) S(1) & S(2) fluxes. T_{rot} is the best fit excitation temperature to several of the lowest rotational transitions. The warm H₂ gas mass (M_{warm}) is computed using the gas temperature listed in any of the preceding three columns. The last column gives the fraction of H₂ gas in the warm component

Object	Type	Ref	Rot. lines observed	T_{01} [K]	T_{12} [K]	T_{rot} [K]	M_{warm} [M_{\odot}]	% of total H ₂
NGC3256	galaxy	a	S0, S1, S2, S5	150 ^a			10 ⁹	3
NGC4038/39	galaxy	b	S1, S2	200 ^a			5.6×10^7	8
NGC4945	galaxy		S0-S3, S5, S7	160	405		8×10^6	1
NGC6946	galaxy	c	S0, S1, S2	179 ^b	380		2.4×10^7	9
Arp220	galaxy	d	S0, S1, S2, S5	150 ^a			1.2×10^6	0.4
Orion Bar	PDR	e	S0-S5, S7, S12, S13, S15, S19, S21	155			–	–
S140	PDR	f	S0-S5, S7, S9, S13	159			–	–
Orion Peak 1	shock	e	S0, S1-S21, S25		578		–	–
Cepheus A West	shock	g	S1-S5, S7, S9			700	–	–
Cepheus A East	shock	h	S0, S1-S8, S9, S10, S11			740	–	–

^a assumed; limits are measured for NGC 3256 (> 140K) and Arp 220 (> 114K)

^b T_{01} recomputed from the original (0-0) S(0) & S(1) fluxes

References: a) Rigopoulou et al. (1996); b) Kunze et al. (1996); c) Valentijn et al. (1996b); d) Sturm et al. (1996); e) D. Rosenthal (priv. comm.); f) Draine & Bertoldi (1999); g) Wright et al. (1996); h) van den Ancker et al. (2000)

- Our mid-infrared ISO spectroscopy does not provide any evidence for the existence of an AGN in the nucleus of NGC 4945. The only high excitation line detected, the 25.9 μm [O IV] line, is most likely produced in shocks associated with the nuclear starburst.
- The AGN, detected in hard X-rays, is unusual in not revealing itself at optical, near-infrared and mid-infrared wavelengths. Hence, either the NLR is extremely obscured ($A_V > 160$), or UV photons from the AGN are absorbed close to the nucleus along all lines of sight, or the AGN is deficient in UV relative to its X-ray flux.
- Many ISM solid state and molecular features have been observed with ISO-PHT-S in the 2.4–11.7 μm range. Most prominent in emission are the PAH features at 3.3, 6.2, 7.7 and 11.2 μm . The strongest absorption features are those of water ice, CO₂ and CO, seen against the nuclear spectrum.

These features show striking similarities to the absorption features seen towards the Galactic center.

- We have studied the physical conditions, excitation and mass of warm H₂, combining IRSPEC and ISO observations of 14 transitions. We derive a visual extinction of 20^{+3}_{-3} mag. to the H₂ emitting region. From the (0-0) S(0) & S(1) lines, we compute a warm (160 K) H₂ gas mass of $2.4 \times 10^7 M_{\odot}$, 9% of the total gas mass inferred from CO. The excitation diagram is best fitted by a power law of the form $dM/dT=4.43 \times 10^{15} T^{-4.793} M_{\odot}/K$. The low excitation temperature of 160 K shows Orion-like shocks not to be representative for the entire emission, and fairly normal PDRs to be perhaps more typical.

Acknowledgements. The authors wish to thank Dietmar Kunze and Fred Lahuis for help in the SWS data reduction and Matt Lehnert, Steve Lord and Eckhard Sturm for stimulating discussions.

References

- Ables J.G., Forster J.R., Manchester R.N., et al., 1987, *MNRAS* 226, 157
- Acosta-Pulido J.A., Klaas U., Laureijs R.J., et al., 1996, *A&A* 315, L121
- Acosta-Pulido J.A., 1999, ISO Explanatory Library, <http://www.iso.vilspa.esa.es/users/expl.lib/PHT/chop-report02.ps.gz>
- Armstrong J.T., Barrett A.H., 1985, *ApJS* 57, 535
- Bergman P., Aalto S., Black J.H., Rydbeck G., 1992, *A&A* 265, 403
- Brock D., Joy M., Lester D.F., et al., 1988, *ApJ* 329, 208
- Chiar J.E., Tielens A.G.G.M., Whittet D.C.B., et al., 2000, *ApJ* in press (astro-ph/0002421)
- Clavel J., Schulz B., Altieri B., et al., 2000, *A&A* in press (astro-ph/0003298)
- De Graauw Th., Haser L.N., Beintema D.A., et al., 1996, *A&A* 315, L49
- Draine B.T., 1989, In: Kaldeich B.H. (ed.) *Infrared Spectroscopy in Astronomy*. ESA-SP 290, p. 93
- Draine B.T., Bertoldi F., 1999, In: Cox P., Kessler M.F. (eds.) *The Universe as seen by ISO*. ESA-SP 427, p. 553
- Elvis M., Wilkes B., McDowell J.C., et al., 1994, *ApJS* 95, 1
- Forbes D.A., Norris R.P., 1998, *MNRAS* 300, 757
- Gabriel C., Acosta-Pulido J., Heinrichsen I., Morris H., Tai W.-M., 1997, In: Hunt G., Payne H.E. (eds.) *Proc. of the ADASS VI conference*, ASP Conf. Ser. 125, p. 108
- Genzel R., Lutz D., Sturm E., et al., 1998, *ApJ* 498, 579
- Gerakines P.A., Schutte W.A., Greenberg J.M., van Dishoeck E.F., 1995, *A&A* 296, 810
- Gerakines P.A., Whittet D.C.B., Ehrenfreund P., et al., 1999, *ApJ* 522, 357
- Greenhill L.J., Moran J.M., Herrnstein J.R., 1997, *ApJ* 481, L23
- Guainazzi M., Matt G., Brandt W.N., et al., 2000, *A&A* in press (astro-ph/0001528)
- Hagen W., Tielens A.G.G.M., 1981, *J. Chem. Phys.* 75, 4198
- Heckman T.M., Armus L., Miley G.K., 1990, *ApJS* 74, 833
- Hesser J.E., Harris H.C., van den Bergh S., et al., 1984, *ApJ* 276, 491
- Iwasawa K., Koyama K., Awaki H., et al., 1993, *ApJ* 409, 155
- Kessler M.F., Steinz J.A., Anderegg M.E., et al., 1996, *A&A* 315, L27
- Koornneef J., 1993, *ApJ* 403, 581
- Koornneef J., Israel P.F., 1996, *New Astronomy* 1, 271
- Kunze D., Rigopoulou D., Lutz D., et al., 1996, *A&A* 315, L101
- Lacy J.H., Baas F., Allamandola L.J., 1984, *ApJ* 276, 533
- Lahuis F., Wieprecht E., Bauer O.H., et al., 1998, In: Albrecht R., Hook R.N., Bushouse H.A. (eds.) *Astronomical Data Analysis Software and Systems VII*, ASP Conf. Ser. 145, p. 224
- Leitherer C., Heckman T.M., 1995, *ApJS* 96, 9
- Lemke D., Klaas U., Abolins J., et al., 1996, *A&A* 315, L64
- Lutz D., 1999, In: Cox P., Kessler M.F. (eds.) *The Universe as seen by ISO*. ESA-SP 427, p. 623
- Lutz D., Feuchtgruber H., Genzel R., et al., 1996, *A&A* 315, L269
- Lutz D., Kunze D., Spoon H.W.W., et al., 1998, *A&A* 333, L75
- Marconi A., Maiolino R., Oliva E., et al., 2000, *A&A* in press (astro-ph/0002244)
- Matt G., Guainazzi M., Maiolino R., et al., 1999, *A&A* 341, L39
- Mattila K., Lehtinen K., Lemke D., 1999, *A&A* 342, 643
- Mauersberger R., Henkel C., Whiteoak J.B., et al., 1996, *A&A* 309, 705
- Moorwood A.F.M., Glass I.S., 1984, *A&A* 135, 281
- Moorwood A.F.M., Oliva E., 1988, *A&A* 203, 278
- Moorwood A.F.M., Oliva E., 1994, *ApJ* 429, 602
- Moorwood A.F.M., van der Werf P.P., Kotilainen J.K., et al., 1996a, *A&A* 308, L1
- Moorwood A.F.M., Lutz D., Oliva E., et al., 1996b, *A&A* 315, L109
- Oliva E., Moorwood A.F.M., Drapatz S., et al., 1999, *A&A* 343, 943
- Ott M., 1995, Ph.D. Thesis, Bonn University
- Quillen A.C., Alonso-Herrero A., Rieke M.J., et al., 1999, *ApJ* 527, 696
- Rice W., Lonsdale C.J., Soifer B.T., et al., 1988, *ApJS* 68, 91
- Rigopoulou D., Lutz D., Genzel R., et al., 1996, *A&A* 315, L125
- Rigopoulou D., Spoon H.W.W., Genzel R., et al., 1999, *AJ* 118, 2625
- Risaliti G., Maiolino R., Salvati M., 1999, *ApJ* 522, 157
- Roche P.F., Aitken D.K., 1985, *MNRAS* 215, 425
- Sadler E.M., Slee O.B., Reynolds J.E., Roy A.L., 1995, *MNRAS* 276, 1373
- Schaeidt S., Morris P.W., Salama A., et al., 1996, *A&A* 315, L55
- Siebenmorgen R., Moorwood A., Freudling W., et al., 1997, *A&A* 325, 450
- Sturm E., Lutz D., Genzel R., et al., 1996, *A&A* 315, L133
- Tayal S.S., Gupta G.P., 1999, *ApJ* 526, 544
- Thornley M.D., Förster Schreiber N.M., Lutz D., et al., 2000, *ApJ* in press (astro-ph/0003334)
- Valentijn E., Feuchtgruber H., Kester D.J.M., et al., 1996a, *A&A* 315, L60
- Valentijn E., Van der Werf P.P., De Graauw Th., et al., 1996b, *A&A* 315, L145
- van den Ancker M.E., Tielens A.G.G.M., Wesselius P.R., 2000, *A&A* in press (astro-ph/0004363)
- van Dishoeck E.F., Helmich F.P., de Graauw Th., et al., 1996, *A&A* 315, L349
- Whiteoak J.B., Dahlem M., Wielebinski R., Harnett J.I., 1990, *A&A* 231, 25
- Wieprecht E., Lahuis F., Bauer O.H., et al., 1998, In: Albrecht R., Hook R.N., Bushouse H.A. (eds.) *Astronomical Data Analysis Software and Systems VII*, ASP Conf. Ser. 145, p. 279
- Wright C.M., Drapatz S., Timmermann R., et al., 1996, *A&A* 315, L301
- Young J.S., Scoville N., 1991, *ARA&A* 29, 581