

Abundances of light elements in metal-poor stars

IV. [Fe/O] and [Fe/Mg] ratios and the history of star formation in the solar neighborhood

R.G. Gratton¹, E. Carretta¹, F. Matteucci^{2,3}, and C. Sneden⁴

¹ Osservatorio Astronomico di Padova, Vicolo dell'Osservatorio 5, 35122 Padova, Italy

² Università di Trieste, Dipartimento di Astronomia, Via G.B. Tiepolo 11, 31413 Trieste, Italy

³ SISSA/ISAS, via Beirut, 34014 Trieste, Italy

⁴ University of Texas at Austin and McDonald Observatory, USA

Received 28 February 2000 / Accepted 17 April 2000

Abstract. The accurate O, Mg and Fe abundances derived in previous papers of this series from a homogeneous reanalysis of high quality data for a large sample of stars are combined with stellar kinematics in order to discuss the history of star formation in the solar neighborhood. We found that the Fe/O and Fe/Mg abundance ratios are roughly constant in the (inner) halo and the thick disk; this means that the timescale of halo collapse was shorter than or of the same order of typical lifetime of progenitors of type Ia SNe (~ 1 Gyr), this conclusion being somewhat relaxed (referring to star formation in the individual fragments) in an accretion model for the Galaxy formation. Both Fe/O and Fe/Mg ratios raised by ~ 0.2 dex while the O/H and Mg/H ratios hold constant during the transition from the thick to thin disk phases, indicating a sudden decrease in star formation in the solar neighbourhood at that epoch. These results are discussed in the framework of current views of Galaxy formation; they fit in a scenario where both dissipational collapse and accretions were active on a quite similar timescale.

Key words: stars: abundances – nuclear reactions, nucleosynthesis, abundances – Galaxy: evolution

1. Introduction

Observations of our and outer galaxies allowed to identify various galactic populations: the halo, the thick disk, the thin disk, and the bulge. A model for the evolution of galaxies should explain the origin and properties of these populations, as well as other basic observations like e.g. the relation of Hubble types with local environment, in a unifying scheme. Current models for galaxy formation broadly divide into two families: those considering a dissipational collapse (Eggen et al. 1962; Larson 1974); and those which consider galaxies as the results of the accretion of individual fragments undergoing (some) independent chemical and dynamical evolution (Toomre & Toomre 1972; Searle & Zinn 1978). The transition between the halo and disk

phases is continuous in smooth *dissipational collapse* models, while disk formation is a secondary mechanism in *accretion* ones. Separation between these two classes of models may be quite artificial: in fact various properties of galaxies, like e.g. the light distribution of ellipticals, are well reproduced by inhomogeneous collapses leading to some kind of violent relaxation (Lynden-Bell 1967); on the other side, simulations based on cosmologies dominated by cold dark matter predict that in high density regions galaxies form hierarchically by merging of smaller subunits, while in low density ones they form more gradually by infall of diffuse matter (Frenk et al. 1985). Within this framework, the mechanisms of formation of our own galaxy (the Milky Way) could be determined by examining fossil remnants of the early phases represented by the old (and often metal-poor) stars. The interpretation of the large amount of data gathered in the last years on dynamics and metallicities (as defined by the most easily observed element, Fe) of field stars is however still controversial, and while e.g. some authors consider the thick disk and the bulge (Gilmore et al. 1989) as distinct galactic components, others (Norris 1993) think they are simply the outer (and oldest) part of the disk and central part of the halo respectively. Scenarios of galactic evolution including a hiatus between the formation of the halo and of a secondary disk (Ostriker & Thuan 1975), that were introduced to justify the rarity of metal-poor stars in the solar neighbourhood (Schmidt 1963), are widely applied e.g. to explain the hot, metal-rich intergalactic gas seen in clusters (Berman & Suchkov 1991); however, up to now the observational basis for this hiatus (based on the age gap between open and globular clusters: Demarque et al. 1992; Carraro et al. 1999; the white dwarf cooling sequence: Wonget et al. 1987; Knox et al. 1999; and the Th/Nd ratio nucleo-chronometer for disk and halo stars: Malaney & Fowler 1989; Cowan et al. 1999) are rather weak and controversial.

Relative abundances of O and Fe in stars of different overall metal abundance provide further constraints to the early evolution of the halo and the formation of the galactic disk (Wheeler et al. 1989). O is the main product of hydrostatic He-burning: hence the ejecta of core-collapse supernovae (SNe)

Send offprint requests to: R.G. Gratton

resulting from the evolution of massive stars, usually identified with type II SNe, are expected to be very rich in O (Woosley & Weaver 1986; Thielemann et al. 1990). On the other side, while a fraction of the Fe presently observed in the interstellar medium was synthesized in massive stars (Thielemann et al. 1990), a large fraction of it was likely produced in explosive burning under degenerate conditions in type Ia SNe (Nomoto et al. 1984). Typical lifetimes of the progenitors of type Ia SNe ($\sim 10^8 \div 10^9$ yr) are much longer than those of the progenitors of type II SNe ($\sim 10^7$ yr), and they are actually longer than, or of the same order of, the free fall time in the Galaxy ($\sim 3 \cdot 10^8$ yr); for these reasons the production of the bulk of Fe is expected to be delayed with respect to that of O (Matteucci & Greggio 1986). A clear break in the run of O abundances with overall metallicity $[\text{Fe}/\text{H}]^1$ should signal the onset of the contribution by type Ia SNe, and the location of this break provides an independent estimate for the timescale of star formation during the early stages of galactic evolution (Matteucci & François 1992; hereinafter MF). It should be added that other α -elements (like Mg, Si, and Ca) are expected to behave similarly to O, although for Si and Ca a small contribution by type Ia SNe is also expected.

In the last years various investigations have been devoted to the study of the run of $[\text{O}/\text{Fe}]$ with $[\text{Fe}/\text{H}]$ in halo and disk stars (Wheeler et al. 1989; King 1994; Nissen & Schuster 1997; Fuhrman 1998, 1999; Israelian et al. 1998; Boesgaard et al. 1999). However, a variety of basic questions still lacks of a clearcut answer. The $[\text{O}/\text{Fe}]$ ratio in the halo and the location of the change of slope in the run $[\text{O}/\text{Fe}]$ vs $[\text{Fe}/\text{H}]$ have been addressed by King (1994), who concluded that this change may occur at any value in the range $-1.7 < [\text{Fe}/\text{H}] < -1.0$, corresponding to timescales for the halo formation between $3 \cdot 10^8$ and $3 \cdot 10^9$ yr (MF); this range is large enough to accommodate both a fast, ordered dissipational collapse (Eggen et al. 1962), or a much slower, accretion scenario (Searle & Zinn 1978). Edvardsson et al. (1993) studied the $[\text{O}/\text{Fe}]$ run in disk stars; they suggested that this ratio is constant for $[\text{Fe}/\text{H}] > -0.2$, and argued that the spread in $[\text{Fe}/\text{H}]$ values at any age is an evidence for infall of metal-poor material. Even less understood is the $[\text{O}/\text{Fe}]$ run at intermediate metallicities, corresponding to the thick disk phase (Gilmore et al. 1989; Nissen & Schuster 1997).

The main concerns in previous investigations on O abundances relate (i) to the paucity of samples of significant sizes studied in a homogeneous way, and then to the possible existence of systematic offsets between different sets of data; and (ii) to the discrepancy between O abundances determined using high excitation permitted and low excitation forbidden lines (the first usually observed in dwarfs, the second in giants). Most of this discrepancy can be removed by adopting higher temperatures in the analysis of dwarfs (King 1993); furthermore, the effects of departures from the Local Thermodynamic Equilibrium (LTE) assumption when considering the formation of high excitation permitted O I lines should also be considered, in or-

der to provide abundances at the level of accuracy required for the present purposes. Recent results based on the OH band at the extreme UV edge of ground-based observations have further complicated this issue, suggesting the presence of a quite strong slope in the $[\text{Fe}/\text{O}]$ run with $[\text{Fe}/\text{H}]$ amongst metal-poor stars (Israelian et al. 1998; Boesgaard et al. 1999)

Both a high temperature scale, and consideration of departures from LTE were included in the new homogeneous determinations of abundances of light elements and Fe for a large sample of stars we are presenting in this series of papers. A new, hopefully improved temperature scale based on IRFM temperatures for population I stars and the new model atmospheres by Kurucz (1995) was obtained in Gratton et al. (1996a, Paper I). An extensive discussion of the effects of departures from the assumption of LTE in line formation in the stellar atmospheres was given in Gratton et al. (1999, Paper II); in that discussion, we exploited an empirical calibration of the poorly known cross sections for collisions with H I atoms drawn from a parallel analysis of the spectra of RR Lyrae variables at minimum light, where non-LTE effects are much larger than in the stars here considered (Clementini et al. 1995). Our final abundances for about 300 stars were presented and discussed in Carretta et al. (2000, Paper III). We found that most discrepancies present in earlier works have been removed, our results showing a high degree of internal consistency, at least for stars with effective temperature $T_{\text{eff}} > 4600$ K, although our results are still not easy to be reconciled with the O abundances from the UV OH bands. However, in this paper we will show that once combined with stellar kinematics and compared with models of galactic chemical evolution, our results allow us to throw new light into some of the above mentioned questions: we find that in the framework of homogeneous models, the collapse of the halo and the formation of the thick disk occurred on a short timescale (a few 10^8 yr), although star formation in these environments likely lasted for $1 \div 2$ Gyr; and that there was a sudden decrease in star formation between the thick and thin disk phases, which are then clearly distinct galactic components. It is worth noticing that a decrease in the star formation rate between the formation of the spheroidal components and the disc in disc dominated galaxies, had already been suggested by Larson (Larson et al. 1976). He suggested that such a decrease in the star formation could be due either to the action of tidal forces inhibiting star formation during the later stages of the collapse or to a two-phase structure of the gas, with dense clouds forming rapidly in a spheroidal component and less dense intercloud gas not forming stars and settling to a disc. We argue that our results fit in a scenario in which both collapse and accretion were important in the formation of the Milky Way, these mechanisms having similar timescales; the relative weights of the two contributions in other galaxies might explain the Hubble sequence.

An early, short presentation of the content of this paper was given as a talk at the conference on Formation of the Galactic Halo (Gratton et al. 1996b). Here we give a more complete presentation of our arguments. In the meanwhile, Fuhrman (1998) reached quite similar conclusions, based on an independent careful analysis of a smaller sample of nearby stars.

¹ In this paper we adopt the standard spectroscopic notation: $[\text{X}] = \log_{10}(\text{X})_{\text{star}} - \log_{10}(\text{X})_{\text{Sun}}$ for any abundance ratio X.




Fig. 1. **a** Run of $[\text{Fe}/\text{O}]$ ratio with $[\text{O}/\text{H}]$ for field stars with $T_{\text{eff}} > 4600$ K. Overimposed lines are the predictions from the models of chemical evolution of Matteucci & François (1992): model 1: thick line; model 2: thin line. Error bar is at top left. **b** The same as panel **a** but with models computed assuming that infall is the sum of two exponentials (“halo” and “disk”), both starting at $t = 0$ but with different decay time; in these models, the star formation rate (SFR) has been suddenly decreased after 1 Gyr (thin line) and 2 Gyr (thick line). **c** The same as panel **b**, but with the “disk” infall starting after 2 Gyr and with a threshold of $7 M_{\odot} \text{pc}^{-2}$ for the SFR; thin line represents predictions of a model where the same SFR has been assumed for both halo and disk phases, while the thick line represents a model where the halo SFR has been increased by an order of magnitude

2. Results

Our results for Fe, O and Mg are displayed in Figs. 1 and 2. To further improve homogeneity, we only plotted data for dwarfs (Edvardsson et al. 1993; Tomkin et al. 1992; Nissen & Edvardsson 1992; Zhao & Magain 1990); all these stars have




Fig. 2. Run of $[\text{Fe}/\text{Mg}]$ ratio with $[\text{Mg}/\text{H}]$ for field stars with $T_{\text{eff}} > 4600$ K. Error bar is at top left

$T_{\text{eff}} > 4600$ K. Inclusion of cooler and/or lower gravity stars in our sample does not change the present discussion. The original data (equivalent widths) used in these papers are very consistent with each other, and were measured on high signal-to-noise, high resolution spectra; typical errors of individual equivalent widths determined from independent estimates for the same stars, are ± 2 mÅ, yielding errors of ± 0.04 dex in abundances derived from individual lines. Possible errors in the atmospheric parameters required in the analysis of individual stars (± 80 K in the effective temperatures T_{eff} ; ± 0.3 dex in the surface gravity; $\pm 0.3 \text{ km s}^{-1}$ in the microturbulent velocity) cause uncertainties of ± 0.08 dex in $[\text{Fe}/\text{H}]$, ± 0.07 dex in $[\text{O}/\text{H}]$, and ± 0.07 dex in $[\text{Fe}/\text{O}]$; these are internal errors. Analogous values for Mg are ± 0.05 dex in $[\text{Mg}/\text{H}]$, and ± 0.06 dex in $[\text{Fe}/\text{Mg}]$. Systematic errors should be small since the present analysis is differential with respect to the Sun. In total, we determined $[\text{O}/\text{Fe}]$ ratios for about 160 stars and $[\text{Mg}/\text{Fe}]$ for 197 stars.

Following Wheeler et al. (1989) we have chosen to use O and Mg as reference elements, because they are almost uniquely produced in massive stars. The scatter of data for individual stars is small and compatible with observational errors for stars with $[\text{O}/\text{H}] > -0.5$ and $[\text{Mg}/\text{H}] > -0.6$, although we cannot exclude a gentle (~ 0.1) slope in the run of $[\text{Fe}/\text{O}]$ with $[\text{O}/\text{H}]$; the scatter we obtain for more metal-poor stars (0.10 dex in both $[\text{Fe}/\text{O}]$ and $[\text{Fe}/\text{Mg}]$ ratios; for O the peculiar N-rich dwarf HD 74000 was omitted) may indicate that errors are larger in this abundance range (perhaps due to the apparent star faintness). However, the composition of the interstellar matter could have been slightly inhomogeneous during the early phases of galactic evolution, as suggested by the large spread in the abundances of n-capture

elements found by McWilliam et al. (1995) for $[\text{Fe}/\text{H}] < -2$ (i.e. $[\text{O}/\text{H}] < -1.5$), and the scatter in the Fe/O ratios amongst halo stars found in the careful analysis by Nissen & Schuster (1997). The upper limit for intrinsic star-to-star variations derived from the spread in our data is 0.07 dex, once observational errors are taken into account.

Fig. 1a indicates that the run of $[\text{Fe}/\text{O}]$ with $[\text{O}/\text{H}]$ is quite flat from $[\text{O}/\text{H}] = -2.2$, $[\text{Fe}/\text{O}] = -0.5$ (these are the most metal-poor stars in our sample) to $[\text{O}/\text{H}] = -0.29$, $[\text{Fe}/\text{O}] = -0.36$, where it is possible to locate (with an uncertainty of about ± 0.1 dex) the change of slope due to the onset of the contribution to nucleosynthesis by the bulk of type Ia SNe. A similar result is provided by Fig. 2 for the run of $[\text{Fe}/\text{Mg}]$ with $[\text{Fe}/\text{H}]$. We remark that the values of $[\text{O}/\text{H}]$ and $[\text{Mg}/\text{H}]$ at which the changes of slope occur is large ($[\text{O}/\text{H}] \sim [\text{Mg}/\text{H}] \sim -0.3$).

A direct comparison with existing galactic evolution model is possible for O, for which the predictions of MF models are available. We overposed on Fig. 1a lines representing the predictions given by models 1 and 2 of MF; these models were computed with an e -folding time of 1 Gyr for the infall (halo collapse), and two different laws of star formation: the times required for $[\text{Fe}/\text{H}]$ to raise at $[\text{Fe}/\text{H}] = -1$ (roughly corresponding to the halo-disk transitions) are $1.5 \cdot 10^9$ and $3 \cdot 10^8$ yr respectively. Both models were arbitrarily scaled to match the $[\text{Fe}/\text{O}]$ ratio for halo stars ($[\text{Fe}/\text{H}] < -1$). These scalings only imply small changes in the adopted yields of Fe from type II SNe, which on turn depend on the cut-off mass for the remnants for core-collapse SNe, an ill-defined quantity at present (Timmes et al. 1995). While these scalings do not affect our conclusions, they help to see the main features we like to point out.

Insofar metal-poor stars are considered ($[\text{Fe}/\text{H}] < -0.5$), the rms values of the residuals of points for individual stars around the lines representing the MF models are 0.158 and 0.127 dex for model 1 and 2 respectively: data for metal-poor stars are then better represented by model 2, which considers the production of Fe from massive stars alone during this phase (the shallow slope of $[\text{Fe}/\text{O}]$ with $[\text{O}/\text{H}]$ in this model is due to the dependence of the O/Fe abundance ratio in the ejecta of type II SNe with progenitor mass); while model 1 is in clear disagreement with observations. The conclusion of this (and other comparisons not shown in Fig. 1), is that the raise of O abundances up to $[\text{O}/\text{H}] = -0.3$ occurred on a timescale which is not much longer than the lifetime of type Ia SNe. This timescale will be better quantified later ².

While model 2 is able to better reproduce observations at low metallicities, it fails in the metal-rich range. In fact, χ^2 -tests show that a linear dependence of $[\text{Fe}/\text{O}]$ on $[\text{O}/\text{H}]$ (roughly sim-

ilar to that given by both model 1 and 2) is not a good representation of the observed $[\text{Fe}/\text{O}]$'s for $[\text{O}/\text{H}] > -0.5$, since the scatter of $[\text{Fe}/\text{O}]$ values for $[\text{O}/\text{H}] < -0.1$ is much larger than observational errors. A very similar result might be obtained for Mg. The simplest interpretation is that the Fe content suddenly increased at $[\text{O}/\text{H}] \sim [\text{Mg}/\text{H}] \sim -0.3$; the implication is that there was a phase in which the production of O and Mg (i.e. the formation of type II SNe) felt down to small values, leaving only the Fe producers in activity. This obviously means a sudden decrease in the formation of massive stars and, since we are moving in the framework of a constant initial mass function (MF), in the star formation *tout court*. We found this same feature when considering Si and Ca abundances rather than O ones (these abundances are not discussed in Paper III, but they display trends similar to those found for O and Mg: Gratton et al., in preparation). The synthesis of these elements is also likely related to massive stars (Thielemann et al. 1990).

The phase of low star formation must have lasted enough to allow the explosion of a large fraction of halo and thick disk type Ia SNe, since $[\text{O}/\text{H}]$ and $[\text{Mg}/\text{H}]$ start increasing again from values of $[\text{Fe}/\text{O}]$ and $[\text{Fe}/\text{Mg}] \sim 0.2$ dex higher than that achieved in the previous phase. The simultaneous increase of both $[\text{O}/\text{H}]$ and $[\text{Fe}/\text{O}]$ in stars with $[\text{O}/\text{H}] > -0.5$, $[\text{O}/\text{Fe}] > -0.25$ seems to require that both type Ia and type II SNe contribute to the chemical enrichment during this phase. The regression line through data in this region is:

$$[\text{Fe}/\text{O}] = (0.34 \pm 0.07)[\text{O}/\text{H}] + (0.04 \pm 0.10), \quad (1)$$

based on 71 stars. However, results for Mg in the metal-rich regime are quite different; the regression line through data in this region is:

$$[\text{Fe}/\text{Mg}] = (-0.055 \pm 0.025)[\text{Mg}/\text{H}] - (0.089 \pm 0.076), \quad (2)$$

based on 164 stars. If real, this result indicates that Mg abundances do not increase at the same rate as O ones, suggesting that the O/Mg ratio in the ejecta of type II SNe is a function metal abundance; this could be understood if severe mass loss reduce production of Mg in massive, metal rich stars.

3. Chemical abundances and dynamics

A further basic step can be done by combining information provided by the ($[\text{O}/\text{H}]$ vs $[\text{Fe}/\text{O}]$) and ($[\text{Mg}/\text{H}]$ vs $[\text{Fe}/\text{Mg}]$) diagrams with those obtained from the kinematics of stars in our sample. A caveat should be done here, since stars analyzed in the present paper were collected from various sources, and the (sometimes not well defined) selection criteria introduce important biases: the distribution of stars with O and Mg abundances in our sample is very different from that obtained from a volume limited sample in the solar neighbourhood, and high velocity stars are likely overrepresented amongst the most metal poor ones. In this section we will use this comparison simply to identify stellar populations defined on chemical grounds with those defined from dynamics.

The basic data for this comparison are the O, Mg and Fe abundances drawn from our analysis, and dynamical data determined by Edvardsson et al. (1993), for stars with $[\text{O}/\text{H}] > -0.5$;

² While this conclusion is based on our analysis, it is not obvious that a different conclusion would be obtained even adopting the O abundances from the UV OH band by Israelian et al. (1998) and Boesgaard et al. (1999). In fact, the quite large slope found by these analyses might be explained by a very fast raise in the metal content of the early gas of the galaxy - so fast that only the most massive stars were able to pollute the early interstellar medium; or by models which consider independent chemical evolution of individual halo fragments, like that proposed by Tsujimoto et al. (1999)




Fig. 3. Run of $[\text{Fe}/\text{O}]$ ratio with $[\text{O}/\text{H}]$ for the stars of Fig. 1 having accurate dynamical parameters. Different symbols mark stars in different areas of the diagram. Compare the distribution of points in this diagram with those of the next figure. Error bar is at bottom right




Fig. 4. The same as Fig. 3, but for Mg rather than for O

for stars more metal-poor than this limit in the Tomkin et al. (1992) and Zhao & Magain (1990) samples, similar data were obtained using parallaxes and proper motion from HIPPARCOS (Perryman et al. 1997), radial velocities from the literature (Carney et al. 1994 whenever possible; else data were taken from the SIMBAD database), and the code for galactic orbit calculations by Aarseth as modified by Carraro & Chiosi (1994). Both these samples only include dwarfs in the solar neighbourhood; the Edvardsson et al. sample is essentially a magnitude-limited sample (although the magnitude limit is a - known - function of metallicity), with no kinematical bias; while Tomkin et al. (1992) and Zhao & Magain (1990) selected high proper motion stars, so that a bias toward high velocity stars is certainly present amongst the most metal-poor stars.

The location of the stars with known dynamical data in the ($[\text{O}/\text{H}]$ vs $[\text{Fe}/\text{O}]$) diagram is shown in Fig. 3, where we plotted with different symbols stars in different regions of the diagram: metal-poor stars ($[\text{O}/\text{H}] < -0.5$: group A); Fe-poor O-rich stars ($[\text{O}/\text{H}] > -0.5$, $[\text{Fe}/\text{O}] < -0.25$: group B); and Fe-rich O-rich stars ($[\text{O}/\text{H}] > -0.5$, $[\text{Fe}/\text{O}] > -0.25$: group C). The analogous diagram for Mg is shown in Fig. 4, but in this case group A are stars with $[\text{Mg}/\text{H}] < -0.6$; group B are stars with $-0.5 < [\text{Mg}/\text{H}] < 0$ and $[\text{Mg}/\text{Fe}] < -0.25$; and group C are the remaining metal rich stars. A star-by-star comparison shows that stars are attributed to the same groups when using O and Mg.

In the four panels of Figs. 5 and 6 $[\text{Fe}/\text{O}]$ and $[\text{Fe}/\text{Mg}]$ ratios for these stars are plotted against the rotational velocity around the galactic center, the orbital eccentricity, the maximum height of the orbit over the galactic plane z_{max} , and the age t respectively. The age values are averages of two independent values:

- those determined by Edvardsson et al. (1993) and Schuster & Nissen (1989) derived using an homogeneous procedure from the $(b - y) - c_1$ diagram, calibrated against age using standard isochrones (*i.e.* not O-enhanced) by Vandenberg & Bell (1985)
- and those derived using absolute magnitudes from HIPPARCOS parallaxes (Perryman et al. 1997), interpolating within the grid of isochrones by Padua group (Girardi et al. 2000); in this case the enhancement of O and the other α -elements was considered by modifying the Fe abundance according to the procedure suggested by Straniero et al. (1997)

The two age scales are somewhat different; they were homogenized to an a common (arbitrary) scale before averaging them.

Absolute ages are affected by various uncertainties related to stellar models, while the relative ranking should be quite reliable, as indicated by the good agreement existing between the two sets of age determinations (see Fig. 7). Since basic data for the two age estimates are independent from each other, internal uncertainties on the ages may be obtained by the r.m.s scatter of the differences; in this way, we estimate that typical error bars in $\log(\text{Age})$ for individual stars are ± 0.06 dex (*i.e.* $\pm 15\%$). The internal scatter we get for group A and B is of this same order, indicating that small if any internal scatter exists for these groups.

The average properties for the three chemical groups are listed in Table 1.

The main conclusions we may draw from Figs. 1-6 and Table 1 are:

1. Though some bias may be present, present data support the identification of group A as the halo, of group B as the thick

Table 1. Average chemical and dynamical parameters for stellar populations in the solar neighbourhood determined from observed stars; for each parameter we give the number of stars used to compute the average values, the values, and the rms scatter of individual points around the mean

Parameter	Halo			Thick disk			Thin disk		
	N	Value	RMS	N	Value	RMS	N	Value	RMS
V_{rot} (kms $^{-1}$)	21	48	100	21	144	52	164	206	24
z_{max} (Kpc)	21	2.1	2.3	21	0.6	1.1	164	0.18	0.20
e	21	0.69	0.28	21	0.38	0.19	164	0.14	0.07
$\log t$ (Gyr)	13	1.16	0.08	21	1.12	0.06	164	0.69	0.23
[Fe/H]	29	-1.68	0.41	21	-0.63	0.15	164	-0.19	0.26
[O/H]	29	-1.23	0.40	19	-0.29	0.12	68	-0.18	0.18
[Fe/O] ^a	28	-0.46	0.10	19	-0.36	0.06	67	-0.01	0.11
[Mg/H]	22	-1.42	0.41	21	-0.29	0.13	164	-0.10	0.24
[Fe/Mg]	22	-0.30	0.10	21	-0.32	0.05	164	-0.08	0.08




Fig. 5a–d. Comparison between the [Fe/O] ratios and physical and dynamical parameters (rotational velocity around the galactic centre V_{rot} : panel a; orbital eccentricity e : panel b; maximum height of orbit above the galactic plane z_{max} : panel c; age t : panel d) for the stars of Fig. 4. Different symbols mark stars with different chemical abundances: open squares are stars with $[\text{O}/\text{H}] < -0.5$; filled circles are stars with $[\text{O}/\text{H}] > -0.5$ and $[\text{O}/\text{Fe}] < -0.25$; open triangles are stars with $[\text{O}/\text{H}] > -0.5$ and $[\text{O}/\text{Fe}] < -0.25$. Note the close correspondance between classification of stellar populations in the solar neighbourhood according to chemical and physical criteria




Fig. 6a–d. Comparison between the [Fe/Mg] ratios and physical and dynamical parameters (rotational velocity around the galactic centre V_{rot} : panel a; orbital eccentricity e : panel b; maximum height of orbit above the galactic plane z_{max} : panel c; age t : panel d) for the stars of Fig. 4. Different symbols mark stars with different chemical abundances: open squares are stars with $[\text{Mg}/\text{H}] < -0.6$; filled circles are stars with $-0.5 < [\text{Mg}/\text{H}] < 0$ and $[\text{Mg}/\text{Fe}] < -0.25$; open triangles are the remaining metal rich stars. Note the close correspondance between classification of stellar populations in the solar neighbourhood according to chemical and physical criteria

disk (see e.g. Robin et al. 1996; Norris 1999; Buser et al. 1999), and of group C as the thin disk.

2. In the framework of homogeneous models for the galactic evolution, the formation of stars in the halo and the thick disk was fast (a few 10^8 yr), i.e. shorter than the typical timescale of evolution for the progenitors of type Ia SNe.
3. Given the lack of a clear selection criterion amongst metal-poor stars, our data cannot be used to draw any conclusion about clear breaks between the halo and thick disk populations; the age difference must be small (i.e. the two populations are virtually coeval), else the [O/Fe] and [Mg/Fe]

ratios of the two populations should be different; the ages derived for both groups are well in excess of 10 Gyr. Anyway, this discontinuity is supported by other studies (see Norris 1993). As noticed by Wyse & Gilmore (1992), the dynamical properties of the thick disk are clearly indicative of a flattened intermediate population supported by rotation (these characteristics are drawn from the Edvardsson et al. sample, which are not affected by kinematical biases); while the halo space distribution is much wider, and the system appears to be supported by velocity dispersion rather than by rotation. In our analysis, this last result is likely enhanced




Fig. 7. Comparison between the ages listed by Edvardsson et al. (1989), and the values determined from the location in the colour-magnitude diagram using absolute magnitudes and colours from HIPPARCOS (Perryman et al. 1997)

by the selection biases in the Carney et al. sample. Further strong supports to the present conclusion that the thick disk is a homogeneous, chemically old population is given by the results by Fuhrmann (1998, 1999), and Nissen & Schuster (1997): these authors found that thick disk stars have low $[\text{Fe}/\text{O}]$ and $[\text{Fe}/\text{Mg}]$ ratios (equal or even lower than those found for halo stars of similar metallicity), with very small intrinsic scatter.

4. There was a sudden decrease in star formation during the transition between the thick and thin disk phases, which are then clearly distinct at least from the chemical point of view. The phase of low star formation must have lasted at least 1 Gyr in order to allow for the explosion of the bulk of type Ia SNe; however, the hiatus was not longer than 3 Gyr, else there would be obvious break in the $[\text{Fe}/\text{O}]$ and $[\text{Fe}/\text{Mg}]$ vs age diagrams. We remark that the age difference between the thick disk and the oldest thin disk stars could have been overestimated in Fig. 5d and 6d, since these ages were derived assuming solar ratios of O and Mg to Fe. Again, results from Fuhrmann (1998, 1999) and Nissen & Schuster (1997) well support this picture.
5. Both the $[\text{Fe}/\text{O}]$ and $[\text{Fe}/\text{Mg}]$ ratios for thin disk stars increase with time: in fact there is a significant correlation of these ratios with stellar ages; the linear regression line are:

$$[\text{Fe}/\text{O}] = -(0.19 \pm 0.06) \log t + (0.12 \pm 0.11), \quad (3)$$

based on 68 stars for O, and:

$$[\text{Fe}/\text{Mg}] = -(0.085 \pm 0.026) \log t - (0.024 \pm 0.075), \quad (4)$$

based on 164 stars for Mg. The Pearson correlation coefficients are 0.35 and 0.25 respectively: the probability of getting such high correlation coefficients by chance are $\ll 0.005$ and $\ll 0.01$. The scatter around the mean lines is compatible with the error bars in ages, $[\text{Fe}/\text{O}]$ and $[\text{Fe}/\text{Mg}]$ values.

4. The gap in $[\text{Fe}/\text{O}]$ distribution

If the raise of $[\text{Fe}/\text{O}]$ at constant $[\text{O}/\text{H}]$ is due to a sudden decrease in star formation during the transition between the thick and thin disk phases, then there should be a corresponding gap at $[\text{Fe}/\text{O}] \sim -0.25$ in the distribution of stars with $[\text{Fe}/\text{O}]$. A cumulative diagram of the distribution of $[\text{Fe}/\text{O}]$ and $[\text{Fe}/\text{Mg}]$ values for stars in the Edvardsson et al. sample (Fig. 8 and 9) indeed supports this inference. In the rest of this section, we will evaluate the statistical significance of this gap for O; similar (though slightly less significant) results were obtained for Mg. To this purpose, we performed tests on the distribution of $[\text{Fe}/\text{O}]$ values in that sample, which consists of the brightest *bona fide* single stars with $5600 < T_{\text{eff}} < 6800$ K, having an absolute magnitude from 0.4 to 2 mag smaller than that of the Zero Age Main Sequence at the same $b - y$ colour, in equally spaced bins in $[\text{Fe}/\text{H}]$. Generally, populations of each bin were kept similar, but the two most metal-poor bins ($[\text{Fe}/\text{H}] < -0.6$) have populations about half those of the others, due to the magnitude limit of the survey. However, the 85 stars for which O abundances are available are nearly uniformly distributed with $[\text{Fe}/\text{H}]$, since we have these data for 42% of the stars with $[\text{Fe}/\text{H}] > -0.6$, and for 79% of the stars more metal-poor than this limit. A χ^2 -test confirms that the distribution of stars having $[\text{Fe}/\text{O}]$ values cannot be distinguished from a uniform distribution in $[\text{Fe}/\text{H}]$; anyway, we repeated the following analysis using both a uniform distribution, and distributions obtained by summing a random gaussian distributed term (representing observational errors) to the observed $[\text{Fe}/\text{H}]$'s. The results are very similar.

We first tested the hypothesis that the $[\text{Fe}/\text{O}]$ values are distributed uniformly, as expected if a linear dependence of $[\text{Fe}/\text{O}]$ on $[\text{Fe}/\text{H}]$ holds. We found that this hypothesis can be rejected at a high level of confidence, using both a χ^2 -test and a series of Monte Carlo simulations. Each Monte Carlo simulation included 10,000 extractions of 85 $[\text{Fe}/\text{H}]$ values (using both early described approaches); individual $[\text{Fe}/\text{O}]$'s were the sum of the values deduced from these $[\text{Fe}/\text{H}]$'s using the $[\text{Fe}/\text{O}]-[\text{Fe}/\text{H}]$ law, and of a random gaussian distributed term representing the spread due to observational errors and to the intrinsic star-to-star variations. The simulations were repeated for three values of the standard deviation for this term: 0.05, 0.07, and 0.10 dex; the intermediate value was deduced from our error analysis, and it is equal to the standard deviation from the mean value for thick disk stars; the last one is consistent with the residuals around the best fit relation for thin disk stars. Most of the deviation from a uniform distribution in $[\text{Fe}/\text{O}]$ is due to the excess of stars with $[\text{Fe}/\text{O}] < -0.3$ (corresponding to the thick disk population), and to the lack of stars with $-0.3 < [\text{Fe}/\text{O}] < -0.15$ (the expected location of the gap). It can be noticed that a quadratic




Fig. 8. Cumulative distribution function of $[O/Fe]$ ratios amongst stars in the Edvardsson et al. sample. Note the possible presence of a gap at $[Fe/O] \sim -0.25$

dependence of $[Fe/O]$ on $[Fe/H]$ gives a better fit to observational data. We then repeated the Monte Carlo simulations with a similar $[Fe/O]-[Fe/H]$ law. We found that the probability that the very low number of stars in a bin of 0.1 dex centered at $[Fe/O] = -0.22$ is due to chance is about 0.024 for a uniform distribution with $[Fe/H]$, and 0.013 for a distribution function equal to the frequency distribution (the exact values depend on the assumed observational errors; the above mentioned values refer to the less significant cases obtained assuming that present $[Fe/O]$ values have errors of 0.10 dex). A good significance (chance probability < 0.05) is achieved for 0.1 dex bins centered over the range $-0.22 < [O/Fe] < -0.25$; this suggests that the gap is broader than 0.1 dex.

We conclude that present available data support the hypothesis that there is a gap in the distribution of $[Fe/O]$ as expected from a sudden decrease in star formation during the transition from the thick to thin disk phases; however we think this test should be repeated using a larger and properly selected sample.

5. Comparison with models of galactic chemical evolution

The overall run of $[Fe/O]^3$ with $[Fe/H]$ is certainly related to the delayed Fe synthesis: however, the interplay between star formation rate, progenitor lifetime, and infall can only be cleared out by detailed modelling of galactic chemical evolution. Unfortunately, various aspects of star formation and evolution are still not well understood, so that these models have several free parameters. Furthermore, a fully appropriate comparison between abundances in metal-poor stars and models of galactic evolution should require models which include consistent and detailed treatment of both chemistry and dynamics. However, this is still beyond current computational capabilities, and at

³ In this section we will only use Fe and O abundances; however, results obtained by considering Mg abundances would be very similar.




Fig. 9. The same as Fig. 7, but for Mg rather than for O. Note again the possible presence of a gap at $[Fe/Mg] \sim -0.25$

present dynamics has to be introduced into chemical evolution models parametrically, increasing even more the number of free parameters. In order to explore allowed ranges for some of the relevant quantities, we then computed a large number of single-zone models (with infall of original unprocessed material) appropriate for the solar neighborhood; the code we used is an improved version of that of MF. The main advantage of this approach is the reduction in the number of free-parameters, although results of our comparisons strictly apply only to homogeneous collapse models, and not e.g. to accretion scenarios. We will see in the next section that even with these limitations, results of our comparisons are enough to suggest that the best scenario required to explain the star formation history in the solar neighborhood should include both dissipational collapse and accretion.

When comparing the predictions of our models with observations, we considered a wide range of constraints, including present abundances in the interstellar medium, current gas density, star formation and SN rates, the distribution of abundances among G-dwarfs, the run of $[Fe/H]$ with age, IMF, etc. (Matteucci 1991), and retained only those models which give an overall match to all of them. We find that in the best cases, rms values of the residuals of $[Fe/O]$'s for individual stars around lines representing our models are ~ 0.10 dex for halo and thick disk stars, and ~ 0.08 dex for thin disk stars, in good agreement with observational errors, as shown by Monte Carlo simulations where both errors in $[Fe/O]$ and $[O/H]$ were taken into account (similar simulations were also used to reject those models providing poor fits). Predictions about the $[Fe/O]$ ratios provided by some of these models are plotted in Fig. 3b and c, overposed to the same observational points shown in Fig. 3a.

On the whole, we found that only models having two distinct infall episodes, the first connected to the halo (and thick disk), and the second to the thin disk, fit observational data (see also Chiappini et al. 1997 for similar models); while models with

a single infall episode do not give a good match even when star formation law was changed with time, due to either large accumulation of gas during the phase corresponding to the raise of the [Fe/O] ratio, or lack of gas at present (depending on the rate of decay). Decay of the halo infall rate should be fast in order to reproduce the almost flat run of [Fe/O] with [O/H] in the halo and thick disk: the e -folding time is shorter than or equal to the adopted time delay for type Ia SNe, i.e. ~ 0.5 Gyr. The halo infall should have contributed no more than 20-30% of the present density in the solar neighborhood (and the fraction in stars should be about half that). The decay of the thin disk infall should be rather slow (≥ 4 Gyr, which implies a present infall rate $\geq 1 M_{\odot} \text{ pc}^{-2} \text{ Gyr}^{-1}$), else there would be not enough gas at present, the present interstellar medium should be too metal rich, the observed roughly linear run of [Fe/O] with [O/H] in the thin disk would not be reproduced, and there would be an excess of moderately metal-poor stars ([Fe/H] ~ -0.5). This value for the present infall rate is larger than the upper limit deduced from observations ($\sim 0.7 M_{\odot} \text{ pc}^{-2} \text{ Gyr}^{-1}$: Mirabel 1989). This inconsistency might be removed by either assuming that star formation law changes with time (the efficiency should then be larger in the halo than in the thin disk), or that the initial mass function (IMF) is flatter at large masses (slope ~ 1.3) than that adopted in most of our models (Scalo 1986) (note that in order to avoid O overproduction with this flatter IMF, an upper limit of $40 \div 50 M_{\odot}$ has to be adopted for type II SNe). Both these alternatives do not contradict basic constraints.

Models where both infall episodes start at high values at the beginning and then decay (Fig. 3b), as well as models where the disk infall is delayed with respect to the halo one (Fig. 3c), fits data quite well. It should be noticed that in the first class of models the star formation must be arbitrarily lowered at the thick-thin disk transition (that should occur ~ 1 Gyr after beginning), and there is no hiatus: the raise of the [Fe/O] ratio at constant [O/H] is due to Fe production by the large number of type Ia SNe from the halo-thick disk, while in the meantime only small amounts of O are produced, due to the decreased star formation rate, which barely compensate for the dilution by metal-poor infalling material (the [O/H] ratio starts increasing again when most gas is consumed and the infall rate decays down to small values). The less appealing aspect of this class of models is the large fraction of metal-poor stars, that only can be reconciled with the small number of metal-poor G-dwarfs observed in the solar neighborhood by assuming that the scale height is a strong function of metallicity. On the other side, the second class of models naturally yield to a hiatus if a threshold (surface) gas density for star formation is adopted (Kennicutt 1989). A threshold gas density might be expected if the processes involved in star formation are self-regulating, and for a low enough surface gas density the feedback mechanisms that regulate the star formation rate break down (Gallagher & Hunter 1984; Elmegreen 1992; Burkert et al. 1992). Observationally, an approximate value of $1 \div 2 \cdot 10^{21} \text{ atoms cm}^{-2} = 5 \div 10 M_{\odot} \text{ pc}^{-2}$ has been proposed for this threshold from observation of irregular and HI-rich spiral galaxies (Skillman 1986; van der Hulst et al. 1987); this threshold value is close to the current gas den-

sity in the solar neighbourhood (Rana & Basu 1991; Kujiken & Gilmore 1989), as expected in a self-regulating mechanism. With the above mentioned infall rates, a star formation rate with threshold produces a hiatus in star formation at the end of the halo and thick disk phase; in our models, star formation in this early phase lasts for 1.5-2.5 Gyr (inversely depending on the adopted threshold density), and for given yields the final [O/H] value depends on the ratio between the halo infall and the threshold density, because the early chemical evolution is essentially that of a closed box (Phillips et al. 1990). In these models, type Ia SNe begin to contribute to nucleosynthesis during the last phases of halo and thick disk evolution, raising the [Fe/O] ratio; this contribution continues during the hiatus, which must be at least as long as time delay of SN Ia (≥ 0.5 Gyr), in order to raise the [Fe/O] by at least 0.13 dex, that we estimate as the lower limit given by observations. However, the upper limit for hiatus duration cannot be determined from the [Fe/O] run due to a saturation effect.

Part of the gas might be lost at the thick-thin disk transition (due e.g. to a galactic wind induced by SN explosions: Silk 1985); if this occurs, this metal-rich gas will be replaced by more metal-poor infalling gas, and the starting value of [O/H] in the thin disk is lower than the maximum achieved during the thick disk evolution. Models where up to 75% of the gas is lost at this epoch fit data quite well, but they predict the existence of some very O-poor, Fe-rich thin disk stars, which are not present in the observed samples⁴. We think that the overposition of the thick and thin disk sequences over some range in [O/H] can be better explained by noting that stars currently in the solar neighborhood likely formed over a range of galactocentric distances (see François & Matteucci 1993), where the halo surface density and the largest [O/H] values achieved during the halo and thick disk evolution were different. Note that this is also the explanation favoured by other authors (e.g. Edvardsson et al. 1993).

6. Dissipational collapse and accretion

What our data tell us about scenarios of Galaxy formation? The fast rate of star formation for the halo and thick disk ($< 10^9$ yr) is consistent with a smooth dissipational collapse (Eggen et al. 1962; Larson 1974). However, in this case the transition from the thick to thin disk phases should be continuous: some heating mechanism causing the observed sudden decrease in star formation during this transition is missing and should then be introduced. Possible candidates are a high SN rate (Silk 1985; Berman & Suchkov 1991), and merging of smaller galaxies with our own; these mechanisms might also be acting simultaneously.

Our tests with models with a single infall episode (but variable star formation rate) suggest that a simple heating as that produced by SNe cannot reproduce the whole spectrum of observations. Also, the discontinuity in specific angular momentum between halo and thick disk suggests that there is not a smooth transition between these two populations (Wyse & Gilmore

⁴ A few stars with such a chemical composition are indeed known, but they have extreme halo kinematics (King 1997; Carney et al. 1997)

1992). On the other side, merging with gas poor satellite(s) having a mass larger than a few hundredths the disk mass may heat (and destroys) any pre-existing stellar thin disk (Quinn et al. 1993; Walker et al. 1996) and create a thick disk, although this result is not obtained in all simulations (see e.g. Huang & Carlberg 1997), so that it seems to depend on the initial conditions as well as on the properties of the satellite: merging of satellites on prograde orbits more likely produce thick disks, while those on retrograde orbits mainly produce disk tilts (Velazquez & White 1999). The mass range is fixed by the amount of kinetic energy to be injected into the disk: merging with a companion of comparable mass would have transformed the Milky Way into an elliptical, while merging with a very small satellite has only minor effects. If the disk or the satellite contained gas (as indicated by chemical evolution models), merging was likely accompanied by a burst in star formation; a possible support in favour of such a burst is the presence of a numerous population of globular clusters likely connected to the thick disk, distinct from the population connected to the halo (Zinn 1985). Perhaps this burst exhausted the existing gas, contributing to the present thick disk or it may have been concentrated in the bulge regions, far from the solar neighbourhood (as suggested by some simulations: Mihos & Hernquist 1994), or finally the high SN rate might have caused a galactic wind (Berman & Suchkov 1991). Anyway, the present thin disk should have formed later by a secondary process (Ostriker & Thuan 1975); models of disk evolution (Kennicutt 1989; Burkert et al. 1992) then indicate that some time would be required before the critical density was reached and star formation started again. Thick disk stars are then likely older than the oldest stars with thin-disk kinematics and a discontinuity would be expected between the thick and thin disk phases. This agrees with our Fig. 5d, which also indicates that a similar large merging could not have occurred during the last 10 Gyrs, although observations of the Sagittarius dwarf galaxy (Ibata et al. 1994), presently merging with the Milky Way, indicate that minor episodes are still occurring. The n-body simulations (Quinn et al. 1993) indicate that if relatively large merging occurred, the present thick disk would be composed of both stars belonging to the merged satellite(s) and to the original disk; the last one should dominate due to its larger mass.

A pure accretion scenario readily explains the lack of appreciable kinematic and chemical gradients in the outer halo (Carney 1993), but it fails to explain the gradients observed in the inner regions. In a smooth dissipational collapse scenario, like that of Eggen et al., the proposed merging episode appears as an *ad hoc* hypothesis. However, more realistic simulations of galaxy formation by dissipational collapse which include dark matter, gas dynamics, star formation and SN feedback (Katz 1992) suggest that many stars form in the cores of dark matter clumps that form during the collapse: even within this scheme then a long-living thin disk could likely form only after the end of an early chaotic phase. The emerging favoured scenario considers then the inhomogeneous dissipational collapse of the protogalaxy with formation of a few secondary fragments (having of the order of several hundredths or even a few tenths of

the total mass) which are accreted later, as proposed by Norris (1994). In this scenario, a significant fraction of the halo is due to accretion of fragments, explaining its low specific angular momentum (Wyse & Gilmore 1992). Our contributions to this scheme is the consideration that the similar [Fe/O] ratios for thick disk and halo stars can best be understood if timescales for both contraction and accretion are ≤ 1 Gyr (although some later accretion of low-mass fragments is possible), and the suggestion that this scheme might naturally produce a discontinuity between the thick and thin disk phases.

Up to now, the proposed scenario refers to the Milky Way. However, it can be easily extended to other galaxies. A strong support to a scenario of spiral formation including both dissipational collapse and accretion is given by the observation that spirals without significant bulges do not have thick disks (van der Kruit & Searle 1981a, 1981b; Morrison et al. 1994, 1997; Fry et al. 1999; Matthews et al. 1999). Then (i) the presence of thick disks and bulges is not an obvious outcome of galactic formation, but rather depend on some mechanism (e.g. accretion) that may or may not be active; and (ii) their origins are likely related (although likely not on an evolutionary sequence, due to the very different specific angular momentum). This last assertion agrees with the low [Fe/Mg] ratios for stars in the bulge of our own Galaxy (McWilliam & Rich 1994), which suggest that the difference between the ages of the bulk of stars in the bulge, halo and thick disk is ≤ 1 Gyr. The presence in the galactic bulge of stars much more metal-rich than the thick disk stars in the solar neighborhood might be understood by assuming either that these stars formed from metal-enriched material in the burst induced by the same merging episode(s) causing the formation of the thick disk (Mihos & Hernquist 1994); or simply by a pre-existing radial metallicity gradient in the early disk, which is predicted by dissipational collapse models (Larson 1974) and should not be cancelled by later merging episode(s) (Quinn et al. 1993).

Within the mixed scenario, significant thick disks and bulges are related to accretion of satellites. Currently available statistics (Zaritsky et al. 1993) indicate that there is about one satellite with $M_B < -15$ per primary, though we notice that the Milky Way has a much larger number of faint companions and the statistics is based on regions where galaxy density is lower than in the Local Group; furthermore, there is an excess of close satellites (separation < 50 kpc) near the minor axis of primaries (Holmberg 1969; Zaritsky et al. 1997), and it has been suggested that the present population of satellites to the Milky Way represents only a small fraction of the original population (see e.g. Klypin et al. 1999). The excess of close satellite near the minor axis may be explained by assuming that close satellites on low inclination prograde orbits have smaller chance to survive, due to dynamical friction; it has been suggested that these *missing* satellites have merged into the primary (Zaritsky & Gonzalez 1999); their perturbation may have created the thick disk, and their gas may have fueled the bulge. Given the small number of satellites for each primary galaxy, a strong stochastic variation from galaxy to galaxy is expected. The present scenario is then coherent to a picture where the entire Hubble sequence from Sc

to ellipticals might be reproduced by assuming an increasing importance of accretion, which should be correlated with the density of galaxies in the local environment (see e.g. Schweizer 2000).

Acknowledgements. We wish to thank Dr G. Carraro and S. Aarseth for having provided a copy of their code for calculation of galactic orbits; and Dr S. Ryan for interesting comments on an early version of this manuscript.

References

- Berman B.G., Suchkov A.A., 1991, *Ap&SS* 184, 169
 Boesgaard A.M., King J.R., Deliyannis C.P., Vogt S., 1999, *AJ* 117, 492
 Burkert A., Truran J.W., Hensler G., 1992, *ApJ* 391, 651
 Buser R., Rong J., Karaali S., 1999, *A&A* 348, 98
 Carney B.W., 1993, In: Smith G.H., Brodie J.P. (eds.) *The Globular Cluster-Galaxy Connection*. ASP Conf. Ser. 48, p. 234
 Carney B.W., Latham D.W., Laird J.B., Aguilar L.A., 1994, *AJ* 107, 2240
 Carney B.W., Wright J.S., Sneden C., et al., 1997, *AJ* 114, 363
 Carraro G., Chiosi C., 1994, *A&A* 288, 751
 Carraro G., Girardi L., Chiosi C., 1999, *MNRAS*, 309, 430
 Carretta E., Gratton R.G., Sneden C., 2000, *A&A* 356, 238 (Paper III)
 Chiappini C., Matteucci F., Gratton R.G., 1997, *ApJ* 477, 765
 Clementini G., Carretta E., Gratton R.G., et al., 1995, *AJ* 110, 2319
 Cowan J.J., Pfeiffer B., Kratz K.-L., et al., 1999, *ApJ* 521, 194
 Demarque P., Green E.M., Guenther D.B., 1992, *AJ* 103, 151
 Edvardsson B., Andersen J., Gustafsson B., et al., 1993, *A&A* 275, 101
 Eggen O.J., Lynden-Bell D., Sandage A.R., 1962, *ApJ* 136, 748
 Elmegreen B.G., 1992, In: Pfenniger D., Bartholdi P. (eds.) *The Galactic Interstellar Medium*. Springer-Verlag, Berlin, p. 157
 François P., Matteucci F., 1993, *A&A* 280, 136
 Frenk C.S., White S.D.M., Efstathiou G., Davis M., 1985, *Nat* 317, 595
 Fry A.M., Morrison H.L., Harding P., Borosin T.A., 1999, *AJ* 118, 1209
 Fuhrmann K., 1998, *A&A* 338, 161
 Fuhrmann K., 1999, *Ap&SS* 265, 265
 Gallagher J.S., Hunter D.M., 1984, *ARA&A* 22, 37
 Gilmore G., Wyse R.F.G., Kuijken C., 1989, *ARA&A* 27, 555
 Girardi L., Bressan A., Bertelli G., Chiosi C., 2000, *A&AS* 141, 371
 Gratton R.G., Carretta E., Castelli F., 1996a, *A&A* 314, 191 (Paper I)
 Gratton R.G., Carretta E., Matteucci F., Sneden C., 1996b, In: Morrison H., Sarajedini A. (eds.) *Formation of the Galactic Halo... Inside and Out*. ASP Conf. Ser. 92, p. 307
 Gratton R.G., Carretta E., Eriksson K., Gustafsson B., 1999, *A&A* 350, 955 (Paper II)
 Holmberg E., 1969, *Ark. Astron.* 5, 305
 Huang S., Carlberg R.G., 1997, *ApJ* 480, 503
 Ibat R.A., Gilmore G., Irwin M.J., 1994, *Nat* 370, 194
 Israelian G., García López R.J., Rebolo R., 1998, *ApJ* 507, 805
 Katz N., 1992, *ApJ* 391, 502
 Kennicutt R.C. Jr, 1989, *ApJ* 344, 685
 King J.R., 1993, *AJ* 106, 1206
 King J.R., 1994, *AJ* 107, 350
 King J.R., 1997, *AJ* 113, 2302
 Klypin A., Kravtsov A.V., Valenzuela O., Prada F., 1999, *ApJ* 522, 82
 Knox R.A., Hawkins M.R.S., Hambly N.C., 1999, *MNRAS* 306, 736
 Kuijken K., Gilmore G., 1989, *MNRAS* 239, 604
 Kurucz R.L., 1995, CD-ROM 13
 Larson R.B., 1974, *MNRAS* 166, 585
 Larson R.B., 1976, *MNRAS* 176, 31
 Lynden-Bell D., 1967, *MNRAS* 136, 101
 Malaney R.A., Fowler W.A., 1989, *MNRAS* 237, 67
 Matteucci F., 1991, In: Lambert D. (ed.) *Frontiers of Stellar Evolution*. PASPCS 20, p. 539
 Matteucci F., François P., 1992, *A&A* 262, L1 (MF)
 Matteucci F., Greggio L., 1986, *A&A* 154, 279
 Matthews L.D., Gallagher J.S., van Dreil W., 1999, *AJ* 1188, 2751
 McWilliam Rich M.J., 1994, *ApJS* 91, 749
 McWilliam A., Preston G.W., Sneden C., Searle L., 1995, *AJ* 109, 2757
 Mihos J.C., Hernquist L., 1994, *ApJ* 425, L13
 Mirabel I.F., 1989, In: Moles M., Tenorio-Tagle G., Melnick J. (eds.) *Structure and Dynamics of the Interstellar Medium*. Springer, Berlin, p. 396
 Morrison H.L., Miller E.D., Harding P., Stinebring D.R., Boroson T.A., 1997, *AJ* 113, 2061
 Morrison H.L., Boroson T.A., Harding P., 1994, *AJ* 108, 1191
 Nissen P.E., Edvardsson B., 1992, *A&A* 261, 255
 Nissen P.E., Schuster W.J., 1997, *A&A* 326, 751
 Nomoto K., Thielemann F.K., Yokoi I., 1984, *ApJ* 286, 644
 Norris J.E., 1993, In: Smith G.H., Brodie J.P. (eds.) *The Globular Cluster-Galaxy Connection*. ASP Conf. Ser. 48, p. 259
 Norris J.E., 1994, *ApJ* 431, 645
 Norris J.E., 1999, *Ap&SS* 265, 213
 Ostriker J.P., Thuan T.X., 1975, *ApJ* 202, 353
 Perryman M.A.C., et al., 1997, *A&A* 323, L49
 Phillipps S., Edmunds M.G., Davies J.I., 1990, *MNRAS* 244, 168
 Quinn P.J., Hernquist L., Fullagar D.P., 1993, *ApJ* 403, 74
 Rana N.C., Basu S., 1991, *A&A* 265, 499
 Robin A.C., Haywood M., Crézè M., Ojha D.K., Bienaymé O., 1996, *A&A* 305, 125
 Scalo J.M., 1986, *Fund. Cosmic Phys.* 11, 1
 Schmidt M., 1963, *ApJ* 137, 758
 Schuster W.J., Nissen P.E., 1989, *A&A* 222, 69
 Schweizer F., 2000, *astro-ph/0002263*
 Searle L., Zinn R., 1978, *ApJ* 225, 357
 Silk J., 1985, *ApJ* 297, 9
 Skillman E.D., 1986, In: Lonsdale Persson C.J. (ed.) *Star Formation in Galaxies*. NASA, Washington, p. 263
 Straniero O., Chieffi A., Limongi M., 1997, *ApJ* 490, 425
 Thielemann F.K., Hashimoto M., Nomoto K., 1990, *ApJ* 349, 222
 Timmes F.X., Woosley S.E., Axelrod T.A., 1995, *ApJS* 98, 617
 Tomkin J., Lemke M., Lambert D.L., Sneden C., 1992, *AJ* 104, 1568
 Toomre A., Toomre J., 1972, *ApJ* 178, 623
 Tsujimoto T., Shigeyama T., Yoshii Y., 1999, *ApJ* 519, 63
 VandenBerg D.A., Bell R.A., 1985, *ApJS* 58, 561
 van der Hulst J.M., Skillman E.D., Kennicutt R.C., Bothun G.D., 1987, *A&A* 177, 63
 van der Kruit P.C., Searle L., 1981a, *A&A* 95, 105
 van der Kruit P.C., Searle L., 1981b, *A&A* 95, 116
 Velazquez H., White S.D.M., 1999, *MNRAS* 304, 254
 Walker I.R., Mihos J.C., Hernquist L., 1996, *ApJ* 460, 121
 Wheeler J.C., Sneden C., Truran J.W., 1989, *ARA&A* 27, 279
 Woosley S.E., Weaver T.A., 1986, In: Mihalas D., Winkler K.A. (eds.) *IAU Coll.* 89, p. 91
 Wonget D.E., Hansen C.J., Liebert J., et al., 1987, *ApJ* 315, L77
 Wyse R.F.G., Gilmore G., 1992, *AJ* 104, 144
 Zaritsky D., Gonzalez A.H., 1999, *PASP* 111, 1508
 Zaritsky D., Smith R., Frenk C., White S.D.M., 1993, *ApJ* 405, 464
 Zaritsky D., Smith R., Frenk C.S., White S.D.M., 1997, *ApJ* 478, L53
 Zinn R., 1985, *ApJ* 293, 424
 Zhao G., Magain P., 1990, *A&AS* 86, 65