

Letter to the Editor

Hard X-ray emission from IC443: evidence for a shocked molecular clump?

F. Bocchino^{1,2} and A.M. Bykov³

¹ Astrophysics Division, Space Science Department of ESA, ESTEC, Postbus 299, 2200 AG Noordwijk, The Netherlands

² Osservatorio Astronomico di Palermo, Piazza del Parlamento 1, 90134 Palermo, Italy

³ A.F. Ioffe Institute for Physics and Technology, St. Petersburg, Russia, 194021

Received 17 August 2000 / Accepted 22 September 2000

Abstract. We report *BeppoSAX* observations of spatially resolved hard X-ray emission from IC 443, a supernova remnant interacting with a molecular cloud. The emission is shown to come from two localized features spatially correlated with bright molecular emission regions. Both hard X-ray features have soft X-ray counterparts, in one case shifted by $\sim 2'$ arcmin toward the remnant interior. The spectra of X-ray emission from both isolated features have photon index $\lesssim 2.0$ in the MECS regime. The emission detected from the remnant with PDS detector extends up to 100 keV. We discuss the observed properties of the hard X-ray features in relation to non-thermal emission from shocked molecular clumps and pulsar wind nebula.

Key words: acceleration of particles – radiation mechanisms: non-thermal – ISM: clouds – ISM: cosmic rays – ISM: individual objects: IC443 – ISM: supernova remnants

1. Introduction

Massive stars that are the likely progenitors of core collapsed supernovae are expected to be spatially correlated with molecular clouds. IC 443 (G189.1+3.0) is one of the best space laboratories to study rich phenomena accompanying supernova interaction with a molecular cloud. This was established from observed shock-excited molecular line emission from OH, CO and H₂ (e.g. Burton et al. 1988; van Dishoeck et al. 1993; Cesarsky et al. 1999). Observations of OH masers and CO clumps towards IC 443, W28 and W44 also provide convincing evidence of molecular cloud interaction in this case (Claussen et al 1997).

IC 443 was a target of X-ray observations with *HEAO 1* (Petre et al. 1988), *Ginga* (Wang et al. 1992), *ROSAT* (Asaoka & Aschenbach 1994) and *ASCA* (Keohane et al. 1997, K97 hereafter). The soft X-ray 0.2–3.1 keV surface brightness map of IC 443 from the *Einstein* Observatory (Petre et al. 1988) shows bright features in the northeastern part of the remnant as well as bright soft emission from the source central part. The

presence of nearly uniform X-ray emission from the central part of the remnant was clearly seen also by *ROSAT* (Asaoka & Aschenbach, 1994), and corresponds to the emission from hot ($T \sim 10^7$ K), low density gas interior to the shock. *ASCA GIS* observations by K97 discovered the localized character of the hard X-ray emission. They concluded that most of the 2–10 keV *GIS* photons came from an isolated emitting feature and from the southeast elongated ridge of hard emission. Preite-Martinez et al. (1999) have reported a hard (> 14 keV) component with *BeppoSAX/PDS* and two hot spots corresponding to the *ASCA* sources of K97 using the MECS.

The supernova remnants (SNRs) that are likely candidates to be γ -ray sources in *CGRO EGRET* observations (Esposito et al. 1996) also show evidence for interaction with molecular gas. In the case of IC 443, the molecular line emission region is partially inside the *EGRET* γ -ray detection circle.

Theoretical models of SNRs interacting with clouds were discussed recently by Sturmer et al. (1997), Baring et al. (1999), Chevalier (1999). The non-thermal multi-wavelength spectrum of a SNR interacting with a molecular cloud was studied by Bykov et al. (2000). They showed that the propagation of a radiative shock wave within a molecular cloud lead to a substantial non-thermal emission both in hard X-rays and in γ -ray. The complex structure of molecular cloud consisting of dense massive clumps embedded in the inter-clump medium could result in localized sources of hard X-ray emission correlated with both bright molecular emission and extended source of non-thermal radio and γ -ray emission. In this Letter, we present the archive *BeppoSAX* observations of IC443 and, in particular, we discuss the features detected in the MECS hard X-ray mosaiced map of the remnant.

2. Observations and data analysis

Results from the the Medium-Energy Concentrator Spectrometer (MECS; 1.8–10 keV; Boella et al. 1997) and the Phoswich Detection System (PDS; 15–300 keV; Frontera et al. 1997) on-board *BeppoSAX* are presented.

Send offprint requests to: byk@astro.ioffe.rssi.ru




Fig. 1. MECS vignnetted and exposure corrected mosaic of the four IC443 observations listed in Table 1. Grayscale is linear and contours are 20%, 40%, 60% and 80% of the peak (3.6 and 2.2×10^{-3} cnt s^{-1} arcmin $^{-1}$ for the soft and hard image, respectively). In the right panel, the sources discussed in the text are marked A, B and C, and dashed contour correspond to a $v = 1 - 0.5(1)$ molecular H line intensity of 6.6×10^{-14} erg cm $^{-2}$ s $^{-1}$ reported by Burton et al. (1988); arrows indicate the location of peaks with an H line intensity at least 14 times the contour levels.

Table 1. Public BeppoSAX Observations of IC443

Obs.	T_{exp}^{MECS} ksec	Coord (J2000)	Date
IC443H	21.5	$6^h 17^m 05^s + 22^d 22^m 19^s$	11/4/1999
IC443E	48.4	$6^h 18^m 00^s + 22^d 34^m 01^s$	11/4/1998
IC443C	67.1	$6^h 16^m 01^s + 22^d 28^m 55^s$	18/10/1997
IC443N	23.8	$6^h 17^m 12^s + 22^d 48^m 50^s$	20/10/1997

We have used all the 4 observation of IC443 available in the public *BeppoSAX* archive; Table 1 lists their coordinates and exposure times. Good data were selected from intervals when the elevation angle above the Earth’s limb was $>4^\circ$ and when the instrument configurations were nominal, using the SAXDAS 2.0.0 data analysis package.

Fig. 1 shows the MECS mosaics of IC443 in different bands. The images are vignetting and exposure corrected. The 2.0-4.0 MECS images show an emission morphology similar to the GIS broad-band maps presented by K97. On the other hand, the hard MECS image above 4 keV is very different. There is weak diffuse extended emission in the north of the remnant, and two strong compact sources, Src A and B. Table 2 reviews the properties of the two sources. In Fig. 1, we also marked the weak extended hard X-ray emission with C. Both the Src A and B were observed by K97; Src A corresponds to the hard source named “HXF”, while Src B corresponds to the hard extended emission called “The Ridge” (also reobserved by Olbert et al. 2000). In the hard X-ray image in Fig. 1 we have also reported the contour of the molecular hydrogen line emissivity observed by Burton et al. (1988), which shows that the hard X-ray sources are closely correlated with it.

Table 2. Compact sources detected in the hard MECS images.

Name	Count-rate ^a (s $^{-1}$)	Position (J2000)
Src A	$3.7 \pm 0.2 \times 10^{-2}$	$6^h 17^m 6.3^s + 22^d 21^m 37^s$
Src B	$5.9 \pm 0.6 \times 10^{-3}$	$6^h 18^m 2.7^s + 22^d 27^m 28^s$

^a In the 4.0-10.0 keV energy range.

In Fig. 2 we report the 4.0-10.5 keV profile of Src A and Src B compared with the MECS PSF profile in the same energy range. The PSF profile was derived from a BeppoSAX observation of the very bright point source Cyg X-1. Src A seems to be extended, while Src B is consistent with the expected PSF. The latter was reported as an elongated ridge by K97, but we note that the ASCA GIS pointing direction is such that the location of Src B is seen at large off-axis angle, which can account for most of source extension observed by ASCA. Instead, the BeppoSAX Src A and B are placed only at $5'$ and $1'$ off-axis, where the instrumental distortion is at its minimum.

We have also investigated if Src A and Src B have soft X-ray counterpart using archive ROSAT observations. Fig. 4 is a PSPC 0.2-2.0 keV image of the region containing MECS Src A and B, with the MECS 4.0-10.5 keV contours of Fig. 1 (right panel) overimposed. Src A has an HRI counterpart quoted by K97 and the PSPC WGACAT source 1WGA J0617.1+2221 with a rate of $2.8 \pm 0.3 \times 10^{-2}$ cnt s $^{-1}$ is located inside the BeppoSAX error circle. Src B has a weaker PSPC counterpart indicated by the arrow in Fig. 4, located $\sim 1.5'$ north of the MECS source centroid and not reported in other PSPC catalogs. We have verified that the extended hard X-ray emission we have labelled with C




Fig. 2. Hard X-ray profile (4.0-10.5 keV) of the MECS Src A and B (dashed line) compared with the expected profile of point-like source (solid line) derived from a long MECS observation of the X-ray binary Cyg X-1.




Fig. 3. MECS and PDS spectrum of Src A. The power law model which best fit the data (Table 3) is also reported with residuals.

in Fig. 1 has a thermal origin and it represents the weak hard tail of the bright soft X-ray nebula visible in the PSPC image.

We have performed spectral fits of the MECS spectra of Src A and B using the MEKAL optically thin thermal plasma model modified by the interstellar absorption assuming the standard abundances. Spectra were extracted from circular regions with radii of 4' and 3' for Src A and B, respectively. We have used the IC443H observation for Src A and IC443E observation for Src B. MECS spectra have been rebinned to have at least 20 counts per channel. The MECS background was collected in an annulus between 5' and 8' for Src A, while we have used the standard background for Src B which is immersed in the thermal emission and a proper background region could not be found around it.

A thermal origin for the spectrum of Src A is strongly rejected ($\chi^2/d.o.f. = 111/41$), while a power law model nicely fit the data (Table 3). The spectrum of Src B cannot be reproduced by a power law or by a thermal emission model alone




Fig. 4. 0.2-2.0 keV PSPC image of the region containing the two MECS hard X-ray sources. The hard X-ray contours of Fig. 1 are overlaid. The arrow shows the soft X-ray counterpart of Src B.

Table 3. Summary of MECS spectral fitting results.

Model	kT keV	γ	χ^2/dof
Src A			
P.L. only	–	$1.96^{+0.21}_{-0.12}$	34/41
Src B			
P.L.+MEKAL	$1.07^{+0.23}_{-0.31}$	$1.5^{+0.9}_{-0.9}$	41/33

($\chi^2/d.o.f. = 61/33$ and $78/33$ respectively) but requires a combination of the two, reported in Table 3.

We have also analyzed the PDS spectra. Background was subtracted using the OFF collimator positions; we have verified that no other contaminating sources are present in the background positions. Spectra were collected using the Variable Rise Time Threshold mode and have been rebinned following a logarithmic scheme suggested by the PDS hardware group to not undersample the instrument spectral resolution. Table 4 reports the observed countrates and flux in different energy ranges. A joint fit of Src A MECS spectrum and IC443H PDS spectrum (shown in Fig. 3) yields best-fit results similar to the ones reported in Table 3 and a PDS/MECS normalization ratio of 1.40 ± 0.45 . This value is consistent with the expected value of 1, also considering that the expected contribution of Src B in the PDS spectrum of IC443 could be up to $\sim 50\%$. As for Src B, the joint fit yields

Table 4. Observed background subtracted count-rates and unabsorbed fluxes in different energy ranges, MECS for $E < 10$ keV and PDS for $E > 10$ keV.

Band keV	Count-rate (10^{-2} s^{-1})	Flux $\times 10^{-12}$ ^(a) erg $\text{cm}^{-2} \text{ s}^{-1}$
Src A		
2–10	8.2 ± 0.2	7.5 ± 1.2
15–30	11.5 ± 3.0	9.1 ± 2.4
30–100	7.2 ± 4.3	11.1 ± 6.6
100–220	5.8 ± 3.7	41.5 ± 26.7
Src B		
2–10	2.0 ± 0.1	1.8 ± 0.2^1
15–30	4.3 ± 2.4	3.4 ± 1.9
30–100	< 3.9	< 1.1

^a Computed using the best-fit models and values reported in Table 3.

a PDS/MECS ratio of 2.8 ± 2.5 , we argued that the PDS spectra of IC443E is heavily contaminated by the brighter Src A. In fact, using the best-fit model of Src A and considering the triangular spatial response of the PDS collimator, we predict a Src A contribution of $\sim 6 \times 10^{-2} \text{ cnt s}^{-1}$ in the PDS spectra of Src B between 15 and 30 keV, consistent with the observed rate (Table 4). Therefore, a proper analysis of the PDS spectra of Src B cannot be done.

3. Discussion

The compact hard X-ray features may be interpreted in terms of the interaction between the remnant shock and a molecular cloud. Bykov et al. (2000) have presented a model of non-thermal emission from an evolved SNR interacting with a molecular cloud, but the predicted diffuse hard X-ray emission in the MECS bandwidth is below the sensitivity threshold. On the other hand, several clumps have been observed in the molecular cloud near IC443. The spatial scales of the molecular clumps from IR line observations are in the range from $150''$ down to $1''$. The non-thermal X-rays from clumps were modelled by Bykov et al. (2000), who predicted a hard spectrum with a photon index below 2.0. The emission is originated from electrons of energy below 1 GeV. The predicted flux density is $\sim 6 \times 10^{-5} \text{ keV cm}^{-2} \text{ s}^{-1} \text{ keV}^{-1}$ at 10 keV for a 30 km s^{-1} shock in a clump of half parsec radius and density of 10^4 cm^{-3} . Using a power law with $\gamma = 1.5$, this corresponds to $\sim 3 \times 10^{-2} \text{ MECS cnt s}^{-1}$ in the 4.0–10.5 keV energy range and $\sim 10^{-2} \text{ PDS cnt s}^{-1}$ in the 15–30 keV. The predicted count-rates are consistent with the observed rate of Src B (Table 4), having in mind that 10^4 cm^{-3} density is some representative number and other parameters could easily account for the difference. Because of hard spectrum and heavy photoelectric absorption, localized spots of a scale of a few arcmin would be seen only in hard X-rays. Thus, the compact Src B correlated with bright spot

of molecular hydrogen emission (Fig. 1) can be shocked molecular clump. The soft X-ray brightening having apparent shift from hard X-ray features towards the SNR interior, observed in the case of Src B (Fig. 4), can be attributed to the effects of the shocked clump edges. The morphology and the spectrum of the Src A are somewhat different from that of Src B.

As already discussed by K97, a low-luminosity pulsar nebula is a plausible explanation for isolated hard X-ray sources (especially for the slightly extended Src A), but we do not expect to see two of them in one SNR². High resolution Chandra or XMM data are needed to distinguish between pulsar nebula and shocked clump interpretation. We have also performed a timing analysis of MECS events collected inside a circle of $3'$ radius in the 4.0–10.5 energy range for both Src A and B, and we found no pulsations at 99% confidence level, with an upper limit of the pulsed fraction of a sinusoidal signal in the $10^{-3} - 64$ Hz frequency range of 5% and 6%, respectively, more stringent than the one derived by K97 using ROSAT data.

We have also estimated the Src A and B EGRET γ -ray $E > 100$ MeV flux contribution, using the best-fit models reported in Table 3, and they are $3.0_{-0.4}^{+0.5} \times 10^{-8}$ and $100_{-92}^{+380} \times 10^{-8}$ photons $\text{cm}^{-2} \text{ s}^{-1}$. The expected γ -ray flux of Src B therefore is compatible with the value of $50.0 \pm 3.0 \times 10^{-8}$ photons $\text{cm}^{-2} \text{ s}^{-1}$ observed by Esposito et al. (1996). Gamma-ray observations with forthcoming *INTEGRAL* mission with expecting angular resolution about $12'$ could better constrain the spectrum of the sources above 30 keV.

Acknowledgements. The authors thank M. Orlandini for his help in analyzing the PDS data and the referee (J.Keohane) for very constructive comments. The work of A.M.B was supported by the INTAS 96-0390 and INTAS-ESA 99-1627 grants. F.B. acknowledges an ESA Research Fellowship.

References

- Asaoka I., & Aschenbach B. 1994, A&A, 284, 573
 Boella G., Chiappetti L., Conti G., et al., 1997, A&AS 122, 327
 Baring M.G., Ellison D.C., Reynolds S.P., et al., 1999, ApJ, 513, 311
 Burton M.G., Geballe T., Brand P., et al., 1988, MNRAS, 231, 617
 Bykov A., Chevalier R., Ellison D., Uvarov, Yu., 2000, ApJ, 538, 203
 Cesarsky D., Cox P., Pineau d. Forets G., et al., 1999, A&A, 348, 945
 Chevalier, R.A. 1999, ApJ, 511, 798
 Claussen M.J., Frail D.A., Goss W.M., et al., 1997, ApJ, 489, 143
 Esposito J.A., Hunter S.D., Kanbach G., et al., 1996, ApJ, 461, 820
 Frontera F., Costa E., Dal Fiume D., et al., 1997, A&AS 122, 371
 Keohane J.W., Petre R., Gotthelf, et al., 1997, ApJ, 484, 350
 Olbert C.M., Clearfield C.R., Williams N.E., Keohane J.W., & Frail D.A., 2000 (in preparation)
 Petre R., Szymkowiak A.E., Seward F.D., et al., 1988, ApJ, 335, 215
 Preite-Martinez A., et al. 1999, AIP Conf. Proc. 510, 73
 Sturmer S.J., Skibo J., Dermer, C. et al. 1997, ApJ, 490, 617
 van Dishoeck E.F., Jansen D., & Phillips T. 1993, A&A 279, 541
 Wang Z.R., Asaoka I., Hayakawa S., et al., 1992, PASJ, 44, 303

² However, Asaoka & Aschenbach (1994) pointed out that the soft X-ray emission is consistent with the presence of two large SNRs, 10^3 and 10^5 yr old.