

## Erratum

## The Galactic disk: study of four low latitude Galactic fields

## A. Vallenari<sup>1</sup>, G. Bertelli<sup>2</sup>, and L. Schmidtobreick<sup>1</sup>

- <sup>1</sup> Padova Astronomical Observatory, vicolo dell'Osservatorio 5, 35122 Padova, Italy
- <sup>2</sup> Department of Astronomy, Padova University, vicolo dell'Osservatorio 5, 35122 Padova, Italy

Astron. Astrophys. 361, 73-84 (2000)

In this paper we correct the following errors introduced in proof:

- 1) In Sect. 6.4, second paragraph, the scale length of the thin disk derived from the data of field F4 is found to be in the range 2–3 kpc (rather than 1.2–1.5 kpc as mistakenly indicated in the text), while values lower than 2 kpc result in a poorer fit to the luminosity functions.
- 2) In the caption of Fig. 11a-d,  $h_r(\text{thick})$  should be replaced with  $h_z(\text{thick})$ . The caption should read:

The luminosity function of Field 4 is displayed with four selected solutions with sech<sup>2</sup> mass distributions:

- a)  $h_r = 3.0 \,\text{kpc}$ ,  $h_z(\text{thin}) = 0.25 \,\text{kpc}$ ,  $h_z(\text{thick}) = 1.5 \,\text{kpc}$ ;
- b)  $h_r = 2.0 \,\text{kpc}$ ,  $h_z(\text{thin}) = 0.28 \,\text{kpc}$ ,  $h_z(\text{thick}) = 1.1 \,\text{kpc}$ ;
- c)  $h_r = 2.5 \text{ kpc}, h_z(\text{thin}) = 0.28 \text{ kpc}, h_z(\text{thick}) = 1.5 \text{ kpc};$
- d)  $h_r = 3.0 \,\text{kpc}, h_z(\text{thin}) = 0.27 \,\text{kpc}, h_z(\text{thick}) = 0.8 \,\text{kpc}$
- 3) In the Conclusions, the scale length of the thin disk found from our data is  $h_r > 2000 \,\mathrm{pc}$  (rather than  $> 1100 \,\mathrm{pc}$ ).

These errors do not affect any of the results or conclusions of the paper.