

Erratum

Line-blanketed model atmospheres for R Coronae Borealis stars and hydrogen-deficient carbon stars

Martin Asplund¹, Bengt Gustafsson¹, Dan Kiselman^{2,3}, and Kjell Eriksson¹

¹ Astronomiska observatoriet, Box 515, S-751 20 Uppsala, Sweden

² NORDITA, Blegdamsvej 17, DK-2100 Copenhagen Ø, Denmark

³ present address: Stockholms observatorium, S-133 36 Saltsjöbaden, Sweden

Astron. Astrophys. **318**, 521–534

Unfortunately the gravities of the models displayed in Fig. 6 and the tickmarks for the optical depths for the comparison model of Schönberner have been mislabeled. The correct figure and caption are shown below. The temperature difference between the two models amounts to about 700 K at the surface ($\log \tau_{\text{Ross}} = -4$) and about 3000 K at depth ($\log \tau_{\text{Ross}} = 1$).




Fig. 6. Comparison of a line-blanketed model atmosphere (solid) with a Schönberger model (dashed) with the same parameters ($T_{\text{eff}} = 7000$, $\log g = 1.0$ [cgs] and C/He = 3%). Rosseland optical depths are marked along the curves

Acknowledgements. We are grateful to Tony Kipper for discovering the mistake.