

Spectroscopic orbit of the ex-eclipsing binary SS Lacertae in the young open cluster NGC 7209

Lina Tomasella¹ and Ulisse Munari²

¹ Osservatorio Astrofisico del Dipartimento di Astronomia, Università di Padova, I-36012 Asiago (VI), Italy

² Osservatorio Astronomico di Padova, Sede di Asiago, I-36012 Asiago (VI), Italy

Received 5 February 1998 / Accepted 24 April 1998

Abstract. The no-longer-eclipsing system SS Lac in the young open cluster NGC 7209 has been recently announced to show a constant radial velocity. Puzzled by this finding, we have monitored the system during 1997 obtaining 24 Echelle+CCD spectra over 8 orbital revolutions. Our spectra reveal a nice orbital motion with periodic splitting and merging of spectral lines from both components.

An accurate orbit has been derived, together with individual masses of the stars. SS Lac presents a moderately eccentric orbit and a probably full synchronization between stellar rotation and orbital revolution.

Key words: binaries: eclipsing – binaries: spectroscopic – stars: individual: SS Lacertae – open clusters: individual: NGC 7209

1. Introduction

According to recent photometry (cf. Vansévičius et al. 1997) SS Lac belongs to the young open cluster NGC 7209. Hoffmeister (1921), Dugan & Wright (1935), Wachmann (1936), Nekrasova (1938) and Kordylewski et al. (1961) described SS Lac as an eclipsing binary of 14.4 days period, with significant eccentricity because the equal depth secondary eclipse occurred at phase 0.57. However, more recent observations by Zakirov & Azimov (1990), Lacy (1992), Mossakovskaya (1993) and Schiller et al. (1996) show that the eclipses no longer occur. The end of the eclipsing phase is set around 1940 by Mossakovskaya (1993) and around 1960 by Lehmann (1991). The latter ascribes the rotation of the plane of the orbit to the presence of an unseen third body in the system, orbiting at a great distance the closer central pair.

Quite puzzling has been the report by Schiller et al. (1991) and Schiller & Milone (1996) that 1983–84 spectra of SS Lac did not reveal the expected double-lined nature of the spectra. Very little, if any, variation in radial velocity was observed. Using available predictions for the rate of change of the SS Lac orbital inclination, Schiller & Milone (1996) expected a semi-amplitude of the radial velocity curve of 150 km sec⁻¹, far in excess of their instrumental resolution. They suggested that

SS Lac could be a triple system suddenly become chaotic or that a close encounter with another NGC 7209 member could have ionized the binary (cf. Schiller 1996).

Stimulated by the Schiller et al. (1991) report, Etzel et al. (1996), Stefanik et al. (1996) and Etzel & Volgenau (1996) announced in IAU Circulars that, according to their preliminary spectroscopic observations, SS Lac is a double-lined system, with indications of variability in the radial velocities. So far no investigation of the orbital motion of SS Lac has appeared in literature.

As an ex-eclipsing binary in a well populated young open cluster, SS Lac clearly deserved further investigations to clarify the whole issue. In this note we report about our spectroscopic monitoring of SS Lac performed to the aim of confirming the binary nature and to derive the spectroscopic orbit.

2. Observations

Twenty-four Echelle+CCD spectra of SS Lac have been obtained with the Asiago 1.82 m telescope during 1997. They cover the range from 4300 to 6600 Å with a resolution of $\lambda/\Delta\lambda \sim 22,500$ at H β (from the FWHM of comparison spectrum Thorium lines). The spectra have been reduced in a standard way with the IRAF software package. A sample of H α profiles recorded on our spectra is shown in Fig. 1, where the periodic splitting and merging of the individual components is evident.

On our spectra we measured only the radial velocities of the H α , H β and H γ lines (the latter over two distinct Echelle orders). A few other metallic lines were indeed visible, but too weak for meaningful measurements. The radial velocities reported in Table 1 come from the hydrogen lines, weighted according to the S/N of the stellar continuum around the line (typically 6:4:3 for H α :H β :H γ , with an average S/N=50–60 around H α).

3. Spectroscopic orbit

Our observations were secured over a time span covering less than 8 revolutions of SS Lac and therefore are not suited for a very accurate period determination. A Deeming-Fourier analysis of data in Table 1 suggests a period of 14.4±0.3 days, very close to the 14.416 days determined from eclipse timing




Fig. 1. A sample of H α profiles of SS Lac reckoned according to the orbital phase. The merging and splitting of the spectral lines is evident. Orbital phases on the left are according to Eq. (2).

by several Authors (e.g. Brancewicz & Dworak 1980). In the computation of the orbit, the period 14.41638 from Eq. (2) has been adopted (see below).

The orbital solution is given in Table 2. With the exception of Ω and eccentricity, the formal errors for the other elements are $\leq 1.5\%$, with a weighted deviation of the observations from the computed orbit of ~ 0.8 km sec $^{-1}$. The solution is graphically presented in Fig. 2.

It is worth to compare the photometric and spectroscopic orbital solutions (cf. Savedoff 1951) through the combination of the eccentricity (e) and longitude of the periastron (Ω) elements (cf. Struve 1948):

$$e \cos \Omega = \frac{2\pi}{P} \frac{(t_2 - t_1 - \frac{1}{2} \times P)}{(1 + \text{cosec}^2 i)} \quad (1)$$

The photometric value is +0.110 (Dugan & Wright 1935), very close to the +0.108 value from the spectroscopic orbital solution in Table 2 (and orbital inclination from Sect. 4.3). Thus the pho-

Table 1. Heliocentric radial velocities of the SS Lac components. Solid line and filled circles in Fig. 2 refer to star a . The quoted errors are probable errors. When the two components were too close in radial velocity for line splitting into individual components, a single value is given (central column). $\text{MJD}_\odot = \text{JD}_\odot - 2450000$.

MJD_\odot	RV_\odot^a	err.	RV_\odot	err.	RV_\odot^b	err.
659.537	14.06	0.21			-49.99	0.14
660.562	34.58	0.39			-68.13	1.07
673.414			-18.69	1.95		
674.337	22.40	1.61			-54.14	1.24
675.361	38.94	1.41			-72.88	2.09
701.448			-20.42	0.90		
714.336	-73.89	1.89			35.91	4.56
714.378	-71.98	1.55			33.63	1.27
715.329			-22.27	1.03		
715.363			-19.00	0.94		
716.315			-17.17	1.08		
716.561			-18.40	0.35		
717.530	20.53	0.28			-58.77	0.15
732.271	25.41	0.38			-65.65	0.74
732.317	25.87	0.09			-67.03	0.26
734.264	44.13	1.01			-83.81	1.57
748.391	41.49	1.04			-85.81	0.80
765.510	26.04	2.00			-70.85	1.19
766.209	0.75	2.10			-60.55	2.00
766.388	0.09	3.45			-47.79	2.38
767.326			-18.14	1.41		
768.345	-64.94	1.09			24.18	1.59
770.223	-102.93	0.50			64.26	0.50
771.205	-97.58	0.50			53.61	0.50

tometric and spectroscopic orbital solutions appear in excellent agreement.

4. Discussion

4.1. Eclipse timing

The $\text{MJD}_\odot=716.315$ spectrum in Fig. 1 presents a perfect radial velocity superposition of the two spectra, thus it corresponds to eclipse conditions. The eclipse ephemeris given by Mossakovskaya (1993)

$$\text{Min } I = 2415900.76 + 14.41619(\pm 0.00013) \times E$$

predicts a +0.456 day shift for the $\text{MJD}_\odot=716.315$ spectrum, which is comparable with the propagation of the uncertainty on the period. Imposing phase coincidence between Mossakovskaya's principal minimum and $\text{MJD}_\odot=716.315$ spectrum, this leads to the improved ephemeris

$$\text{Min } I = 2450716.32(\pm 0.15) + 14.41638(\pm 0.00010) \times E(2)$$

The slightly longer orbital period nearly coincides with those given by Brancewicz & Dworak (1980, $P=14.4163$ days) and by Dugan & Wright (1935, $P=14.41629$ days).

Table 2. Orbital elements for SS Lac. Where appropriate, the second value correspond to the component *b* represented by a dotted line and open circles in Fig. 2. The quoted errors are the formal errors of the orbital solution. The entry “deviation” is the mean weighted deviation of the observed radial velocities from the computed orbital solution (weight = err^{-2} in Table 1). $\text{MJD}_{\odot} = \text{JD}_{\odot} - 2450000$. The error on the masses is $\sim 0.1 M_{\odot}$.

		star <i>a</i>	star <i>b</i>
period	(days)	14.41638	
baricentric velocity	(km sec^{-1})	-21.2 ± 0.3	
semi-amplitude	(km sec^{-1})	74.7 ± 1.1	77.6 ± 1.4
eccentricity		0.122 ± 0.019	
$a_i \sin i$	(AU)	0.098 ± 0.001	0.102 ± 0.002
T_{\odot}	(MJD_{\odot})	741.4 ± 0.2	
Ω	(deg)	332 ± 9	
deviation	(km sec^{-1})	0.87	0.70
<hr/>			
inclination	(deg)	78 ± 5	
mass	(M_{\odot})	2.80	2.69

4.2. Spectral classification

The absence of He I absorptions, the paucity of metallic lines and the very strong Stark broadening of the wings of the Balmer lines suggest a spectral classification around A2 V, virtually identical for the two members (cf. Fig. 1). A more refined classification however needs devoted observations at the head of the Balmer series. Component *b* appears slightly less luminous than component *a*, as suggested by the relative depths of the central Doppler cores and contribution to the overall $H\alpha$ line wings in Fig. 1. A lower brightness agrees with the slightly smaller mass of component *b* (cf. the semi-amplitudes and semi-major axes in Table 2) and the just slightly reduced amplitude of *b* eclipses compared to *a* eclipses (a few hundredth of a magnitude, cf. Dugan & Wright 1935, Mossakovskaya 1993).

4.3. Orbital inclination

The spectral classification of the two components can be turned into a physical radius of $R = 2.25 R_{\odot}$ from interpolation of tabular values given by Allen (1973). The orbital separation from Table 2 is a $\sin i = 0.200 \pm 0.002 \text{ AU} = 42.5 \pm 0.5 R_{\odot}$. Lehmann (1991) has stated that the system presented total eclipses (central eclipses given the nearly identical size of the components) in 1900-1915, and afterward the eclipses monotonically reduced in depth until they disappeared around 1960. From separation and size of the two components it may be computed that at the end of the eclipsing season the inclination was $i \sim 83^{\circ}$. The corresponding linear change in orbital inclination is

$$\frac{di}{dt} = 0.13 \pm 0.01 \quad \text{deg yr}^{-1} \quad (3)$$

which is significantly lower than the $di/dt = 0.18 \text{ deg yr}^{-1}$ obtained by Lehmann (1991) on the assumption of a B7 V classi-




Fig. 2. Orbital solution for SS Lac. The curves give the orbital solution of Table 2. Solid curve and filled circles refer to component *a* in Tables 1 and 2. The crosses mark spectra for which the too small velocity separation caused merging of the two components into a single line. The dashed line is the baricentric velocity ($-21.2 \text{ km sec}^{-1}$). Orbital phase according to Eq. (2).

fication for both components (clearly ruled out by our spectra that show no evidence for HeI absorption lines). From Eq. (3) and the occurrence of central eclipses at the beginning of this century, the current orbital inclination is

$$i_{(1998)} \sim 78^{\circ} \quad (4)$$

The above results indicate that the non-eclipsing season of SS Lac lasts for about 1275 ± 90 years.

4.4. Rotational velocities

The rotational velocity of the components of SS Lac has been derived by comparison with standard stars from Sletteback (1975), which were observed under the same instrumental conditions. The velocity turned out to be the same for both components:

$$V_{rot} = 10 \pm 5 \quad \text{km sec}^{-1} \quad (5)$$

The dependence on inclination (see Eq[4]) has been removed under the assumption that the rotational and orbital axes are parallel. A $V_{rot} = 10 \pm 5 \text{ km sec}^{-1}$ means that rotational and orbital motion are synchronized or close to, the orbital velocity amounting to $V_{orb} = 8 \text{ km sec}^{-1}$.

It is worth noting that SS Lac shows evidence for synchronization when the orbital circularization is still far from being achieved. The two phenomena seems therefore to evolve over quite different time scales, in agreement with the theoretical expectations (cf. Zahn 1977).

Table 3. Heliocentric radial velocity of the weak component of the Balmer lines described in Sect. 4.7. The errors may be quantified as $\pm 1 \text{ km sec}^{-1}$ when $N=3$, ± 2 for $N=2$ and ± 4 for $N=1$. $\text{MJD}_{\odot} = \text{JD}_{\odot} - 2450000$. N = number of Balmer lines used.

MJD	RV_{\odot}	N	MJD	RV_{\odot}	N	MJD	RV_{\odot}	N
	(km/sec)			(km/sec)			(km/sec)	
659.537	-16	1	732.317	-27	1	768.345	-7	1
660.562	-22	3	734.264	-19	3	770.223	-41	1
675.362	-29	1	748.391	-23	2	771.205	-19	1
714.378	-20	1						

The expected synchronization time for a binary with the SS Lac parameters may be estimated from the Tassoul's (1987) formalism to be:

$$t_{syn}(yrs) \geq 2500 \times [P(days)]^{\frac{1}{4}} \quad (6)$$

$$\geq 4 \text{ million years}$$

which is relatively short compared to the NGC 7209 cluster age (300 million yrs according to Lynga 1985). The synchronization status cannot therefore be used to decide if SS Lac is or not a primordial binary of NGC 7209.

4.5. Individual masses

A straightforward application of the Kepler's III law to Table 2 data gives 2.69 and 2.80 M_{\odot} for the masses of the two SS Lac components. The tabular mass from Allen (1973) for the A2 V spectral type is 2.78 M_{\odot} , in excellent agreement.

4.6. Mass-Luminosity relation

Vansevicius et al. (1997) have estimated a 1.0 kpc distance and a $A_V=0.54$ mag extinction to NGC 7209. Using a bolometric correction B.C.= -0.29 mag for the spectral type A2 V (Allen 1973), the bolometric magnitude of the SS Lac components is

$$M_{bol} = +0.25 \text{ mag} \quad (7)$$

corresponding to a luminosity $L = 65 L_{\odot}$. The mean mass for the two components is 2.745 M_{\odot} . At this mass the Mass-Luminosity relation has the numerical form (cf. Bowers & Deeming 1984):

$$\log \frac{L}{L_{\odot}} = 0.479 + 2.91 \log \frac{M}{M_{\odot}} \quad (8)$$

which predicts $L \sim 57 L_{\odot}$ for the SS Lac components. The agreement between the observed and predicted luminosity is satisfactory in view of the uncertainties in the distance, reddening and the approximate bolometric correction.

4.7. The unseen third body

Lehmann (1991) ascribes the rotation of the plane of the orbit to the presence of an unseen third body in the system, orbiting at a great distance the closer central pair. A discussion of this hypothesis is beyond the scopes of the present paper. We may however point out that in our spectra when the Balmer lines show a clear split and the S/N ratio is high, a weak absorption is visible. The $H\alpha$ component does not correspond to known telluric absorptions in this region, and the coincidence between the various Balmer lines favour an interpretation as a *real* stellar line. To the benefit of future Investigators of SS Lac we report in Table 3 the heliocentric velocity of this weak component as measured on our spectra (same weighting as described in Sect. 2), that could in some way be related to the unseen third body.

Acknowledgements. We would like to thank the referee, E.F.Milone, for accurate reading of the manuscript and scrutiny of the data. We thank also M.Rejkuba who secured a few of the spectra used in this investigation and C.Barbieri, G.Cremonese and S.Verani who allowed us to secure a few other spectra during their observing time. R.Margoni contributed useful insights in the statistical treatment of the data.

References

- Allen C.W., 1973, *Astrophysical Quantities*, The Athlone Press
 Bowers R., Deeming T., 1984 *Astrophysics*, Jones and Bartlett Publishers, Boston
 Brancewicz H.K., Dworak T.Z., 1980, *AcA* 30, 501
 Dugan R.S., Wright F.W., 1935, *AJ* 44, 150
 Etzel P.B., Volgenau N.H., Nguyen Q.T., 1996, *IAU Circular No.* 6429
 Etzel P.B., Volgenau N.H., 1996, *IAU Circular No.* 6434
 Hoffmeister C., 1921, *Astron. Nach.* 214, 8
 Kordylewski K., Pagaczewski J., Scafraniec R., 1961, *AcAS* 4, 487
 Lacy C.H., 1992, *AJ* 104, 801
 Lehmann T., 1991, *IBVS* 3610
 Lynga G., 1985, *IAU Symp.* 106, pag. 143
 Mossakovskaya L.V., 1993, *AZh Pis'ma* 19, 87
 Nekrasova S., 1938, *Perem. Zvezd.* 5, 182
 Savedoff M.P., 1951, *AJ* 56, 1
 Schiller S.J., Milone E.F., Zakirov M.M., Azimov A.A., 1991, *BAAS* 23, 879
 Schiller S.J., Milone E.F., 1996, In: Milone E.F., Mermilliod J.C. (eds.) *The Origins, Evolution and destinies of binary stars in Clusters.* ASP Conference Series, Vol. 90, p. 123
 Schiller S.J., Bridges D., Clifton T., 1996, In: Milone E.F., Mermilliod J.C. (eds.) *The Origins, Evolution and destinies of binary stars in Clusters.* ASP Conference Series, Vol. 90, p. 141
 Schiller S.J., 1996, *Astronomy* 24, 46
 Sletteback A., Collins II G.W., Boyce P.B., White N.M., Parkinson T.D., 1975, *ApJS* 29, 137
 Stefanik R.P., Torres G., Caruso J.R., et al., 1996, *IAU Circular No.* 6433
 Struve O., 1948, *PASP* 60, 160
 Tassoul J.L., 1987, *ApJ* 322, 856
 Vansevicius V., Platais I., Paupers O., et al., 1997, *MNRAS* 285, 871
 Wachmann A., 1936, *Astron. Nach.* 258, 361
 Zahn J.P., 1977, *A&A* 57, 383
 Zakirov M.M., Azimov A.A., 1990, *IBVS* 3487